

**Exercise-set 5.**  
**Solutions**

1. Construct a graph  $G$ :  $V(G) =$  people, and the edges are the acquaintances. Then  $\deg(v) \geq 6 = 12/2 \implies$  by Dirac's theorem  $\exists$  a Hamilton cycle.
2. The condition in Ore's theorem holds for  $G \implies \exists$  a Hamilton cycle.
3. Construct a graph  $G$ :  $V(G) =$  people, and the edges are the acquaintances.  $G$  is  $k$ -regular for some  $k$ . If  $k \geq 10 \implies G$  contains a Hamilton cycle, if  $k \leq 9 \implies \overline{G}$  contains a Hamilton cycle.
4. A cycle on  $n$  vertices is like that (check).
5. E.g.  $K_7$  with the edge  $\{u, v\}$  missing and the 8th vertex is connected to  $u$ .
6. Add a new vertex to  $G$ , and connect it to all the old vertices. Then the new graph contains a Hamilton cycle from which we can get a Hamilton path of  $G$ .
7. Add two new non-adjacent vertices to  $G$ , and connect them to all the old vertices. Then the new graph contains a Hamilton cycle from which we can get a Hamilton path of  $G$ .
8. Delete  $v$  from  $G$ . Then the new graph contains a Hamilton cycle from which we can get a Hamilton path of  $G$ .
9. The 8 edges must have pairwise no common endpoints (i.e. be independent). Every second edge of a Hamilton cycle will do (which exists because  $\deg(v) \geq n/2 \forall v$ ).
10. We can add the edges of a Hamilton cycle of  $\overline{G}$ .
11. Need to add  $k$  pairwise non-adjacent edges (from  $\overline{G}$ ).  $\overline{G}$  contains a Hamilton cycle ( $\deg_{\overline{G}}(v) = k, \forall v \in V(G)$ ). Every second edge of it will do.
12. Construct a graph  $G$ :  $V(G) =$  four-element subsets, and they are adjacent, if they have at least two elements in common. Then  $G$  contains a Hamilton cycle.
13. The first graph is not bipartite (contains 5-cycles), but the second graph is.
14. Deleting 2 edges are enough, but less is not, since  $\exists$  2 edge-disjoint odd cycles in  $G$ .
15. The graph determined by the knights and attacks is bipartite (the two classes are to the white and black squares), and each of its degrees is at least 2  $\implies \exists$  a degree  $\geq 3$ .
16. Yes (the two classes of vertices are sequences with an even or odd number of 1's, resp.).
17. No (the complement contains a triangle).
18. The vertices cannot be divided into two classes (degrees).
19. Complete bipartite graphs are like that.
20. The graphs are exactly the odd cycles (so in particular  $n$  must be odd).  $G$  must contain an odd cycle (otherwise  $\chi(G) = 2$ ), and cannot contain more vertices or edges.
21.  $\omega(G) = 3 \implies \chi(G) \geq 3$ , and  $G$  can be colored with 3 colors  $\implies \chi(G) \leq 3$ .
22.  $\omega(G) = 8$  (each row and column is a clique)  $\implies \chi(G) \geq 8$ , and  $G$  can be colored with 8 colors (colors are diagonal)  $\implies \chi(G) \leq 8$ .
23.  $G$  is bipartite (the two classes of vertices are the even and odd numbers, resp.)  $\implies \chi(G) = 2$
24.  $\omega(G) = 4 \implies \chi(G) \geq 4$ , and  $G$  can be colored with 4 colors  $\implies \chi(G) \leq 4$ .
25. a), b)  $\omega(G) = 3 \implies \chi(G) \geq 3$ , but  $G$  cannot be colored with 3 colors (proof!)  $\implies \chi(G) \geq 4$ .  $G$  can be colored with 4 colors  $\implies \chi(G) \leq 4$ .
26.  $\omega(G) = 3 \implies \chi(G) \geq 3$ , but  $G$  cannot be colored with 3 colors (proof!)  $\implies \chi(G) \geq 4$ .  $G$  can be colored with 4 colors  $\implies \chi(G) \leq 4$ .
27.  $\chi(G) \geq \lceil n/2 \rceil$  (at most 2 vertices can get the same color), and  $G$  can be colored with this many colors  $\implies \chi(G) = \lceil n/2 \rceil$ .
28.  $\omega(G) = 10$  (any 10 consecutive numbers form a clique)  $\implies \chi(G) \geq 10$ , and  $G$  can be colored with 10 colors (periodically)  $\implies \chi(G) \leq 10$ .

29.  $\omega(G) = 5$  ( $\{1, 8, 15, 22, 29\}$  is a clique)  $\implies \chi(G) \geq 5$ , and  $G$  can be colored with 5 colors  $\implies \chi(G) \leq 5$ .
30.  $\omega(G) = 11$  ( $\{10, 11, \dots, 20\}$  is a clique)  $\implies \chi(G) \geq 11$ , and  $G$  can be colored with 11 colors  $\implies \chi(G) \leq 11$ .
31.  $\omega(G) = 4$  (the powers of 2 form a clique)  $\implies \chi(G) \geq 4$ , and  $G$  can be colored with 4 colors (using the same color between consecutive powers of 2)  $\implies \chi(G) \leq 4$ .
32.  $\omega(G) = 11$  (prime numbers and 1 form a clique)  $\implies \chi(G) \geq 11$ , and  $G$  can be colored with 11 colors  $\implies \chi(G) \leq 11$ .
33.  $\omega(G) = 5 \implies \chi(G) \geq 5$ , and  $G$  can be colored with 5 colors  $\implies \chi(G) \leq 5$ .
34.  $G$  is  $K_{10}$  with a perfect matching deleted.  $\omega(G) = 5 \implies \chi(G) \geq 5$ , and  $G$  can be colored with 5 colors  $\implies \chi(G) \leq 5$ .
35. Use the greedy coloring in the original (increasing) order of the vertices.
36. Order the vertices: first the exceptional ones, then the rest, and use the greedy coloring.
37. Use the greedy coloring in the decreasing order of the degrees.