

Known Algorithms on Graphs of Bounded Treewidth are Probably Optimal*

Daniel Lokshtanov[†] Dániel Marx[‡] Saket Saurabh[§]

January 14, 2018

Abstract

We obtain a number of lower bounds on the running time of algorithms solving problems on graphs of bounded treewidth. We prove the results under the Strong Exponential Time Hypothesis of Impagliazzo and Paturi. In particular, assuming that n -variable m -clause SAT cannot be solved in time $(2 - \epsilon)^n m^{\mathcal{O}(1)}$, we show that for any $\epsilon > 0$;

- INDEPENDENT SET cannot be solved in time $(2 - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$,
- DOMINATING SET cannot be solved in time $(3 - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$,
- MAX CUT cannot be solved in time $(2 - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$,
- ODD CYCLE TRANSVERSAL cannot be solved in time $(3 - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$,
- For any fixed $q \geq 3$, q -COLORING cannot be solved in time $(q - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$,
- PARTITION INTO TRIANGLES cannot be solved in time $(2 - \epsilon)^{\text{tw}(G)} |V(G)|^{\mathcal{O}(1)}$.

Our lower bounds match the running times for the best known algorithms for the problems, up to the ϵ in the base.

1 Introduction

It is well-known that many NP-hard graph problems can be solved efficiently if the *treewidth* ($\text{tw}(G)$) of the input graph G is bounded. For an example, an expository algorithm to solve VERTEX COVER and INDEPENDENT SET running in time $\mathcal{O}^*(4^{\text{tw}(G)})$ is described in the algorithms textbook by Kleinberg and Tardos [35] (the \mathcal{O}^* notation suppresses factors polynomial in the input size), while the book of Niedermeier [43] on fixed-parameter algorithms presents an algorithm with running time $\mathcal{O}^*(2^{\text{tw}(G)})$. Similar algorithms, with running times on the form $\mathcal{O}^*(c^{\text{tw}(G)})$ for a constant c , are known for many other graph problems such as DOMINATING SET, q -COLORING and ODD CYCLE TRANSVERSAL [5, 19, 26, 27, 50]. Algorithms for graph problems on bounded treewidth graphs have found many uses as subroutines in approximation algorithms [7, 24, 25, 36], parameterized algorithms [4, 21, 23, 34, 41, 49], and exact algorithms [28, 46, 42].

In this paper, we show that any improvement over the currently best known algorithms for a number of well-studied problems on graphs of bounded treewidth would yield a faster algorithm for SAT. In particular, we show if there exists an $\epsilon > 0$ such that

- INDEPENDENT SET can be solved in time $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$, or

*A preliminary version of this paper appeared in the proceedings of SODA 2011 [37].

[†]Department of Informatics, University of Bergen, Norway. daniello@ii.uib.no

[‡]Institute for Computer Science and Control, Hungarian Academy of Sciences (MTA SZTAKI), Budapest, Hungary. dmarx@cs.bme.hu

[§]The Institute of Mathematical Sciences, India. saket@imsc.res.in

- DOMINATING SET can be solved in time $\mathcal{O}^*((3 - \epsilon)^{\text{tw}(G)})$, or
- MAX CUT can be solved in time $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$, or
- ODD CYCLE TRANSVERSAL can be solved in time $\mathcal{O}^*((3 - \epsilon)^{\text{tw}(G)})$, or
- there is a fixed $q \geq 3$ such that q -COLORING can be solved in time $\mathcal{O}^*((q - \epsilon)^{\text{tw}(G)})$, or
- PARTITION INTO TRIANGLES can be solved in time $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$,

then n -variable SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$. Such an algorithm would violate the *Strong Exponential Time Hypothesis* (SETH) of Impagliazzo and Paturi [31]. Thus, assuming SETH, the known algorithms for the mentioned problems on graphs of bounded treewidth are essentially the best possible.

To show our results we give polynomial time many-one reductions that transform n -variable boolean formulas ϕ to instances of the problems in question. Such reductions are well-known, but for our results we need to carefully control the treewidth of the graphs that our reductions output. A typical reduction creates n gadgets corresponding to the n variables; each gadget has a small constant number of vertices. In most cases, this implies that the treewidth can be bounded by $O(n)$. However, to prove the a lower bound of the form $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$, we need that the treewidth of the constructed graph is $(1 + o(1))n$. Thus we can afford to increase the treewidth by at most one per variable. For lower bounds above $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$, we need even more economical constructions. To understand the difficulty, consider the DOMINATING SET problem, here we want to say that if DOMINATING SET admits an algorithm with running time $\mathcal{O}^*((3 - \epsilon)^{\text{tw}(G)}) = \mathcal{O}^*(2^{\log(3 - \epsilon)\text{tw}(G)})$ for some $\epsilon > 0$, then we can solve SAT on input formulas with n -variables in time $\mathcal{O}^*((2 - \delta)^n)$ for some $\delta > 0$. Therefore by naïvely equating the exponent in the previous sentence we get that we need to construct an instance for DOMINATING SET whose treewidth is essentially $\frac{n}{\log 3}$. In other words, each variable should increase treewidth by *less than one*. The main challenge in our reductions is to squeeze out as many combinatorial possibilities per increase of treewidth as possible. In order to control the treewidth of the graphs we construct, we upper bound the *pathwidth* ($\text{pw}(G)$) of the constructed instances and use the fact that for any graph G , $\text{tw}(G) \leq \text{pw}(G)$. Thus all of our lower bounds also hold for problems on graphs of bounded pathwidth.

Complexity Assumption: The *Exponential Time Hypothesis* (ETH) and its strong variant (SETH) are conjectures about the exponential time complexity of k -SAT. The k -SAT problem is a restriction of SAT, where every clause in input boolean formula ϕ has at most k literals. Let $s_k = \inf\{\delta : k\text{-SAT can be solved in } 2^{\delta n} \text{ time}\}$. The Exponential Time Hypothesis conjectured by Impagliazzo, Paturi and Zane [32] is that $s_3 > 0$. In [32] it is shown that ETH is robust, that is $s_3 > 0$ if and only if there is a $k \geq 3$ such that $s_k > 0$. In the same year it was shown that assuming ETH the sequence $\{s_k\}$ increases infinitely often [31]. Since SAT has a $\mathcal{O}^*(2^n)$ time algorithm, $\{s_k\}$ is bounded by above by one, and Impagliazzo and Paturi [31] conjecture that 1 is indeed the limit of this sequence. In a subsequent paper [12], this conjecture is coined as SETH.

ETH is now a widely believed assumption, and has been used as a starting point to prove running time lower bounds for numerous problems [9, 15, 14, 22, 18, 29, 38, 39, 40]. At the time of the conference version of this paper, SETH was largely untouched [44, 11, 16, 20, 21, 17]. The reason for this was two-fold. First, the assumption that $\lim_{k \rightarrow \infty} s_k = 1$ is a very strong one. Second, when proving lower bounds under ETH we can utilize the *Sparsification Lemma* [32] which allows us to reduce from instances of 3-SAT where the number of clauses is linear in the number of variables. Such a tool does not exist for SETH, and this seems to be a major obstruction for showing running time lower bounds for interesting problems under SETH. We overcome this obstruction by circumventing it – in order to show running time lower bounds for algorithms on bounded treewidth graphs sparsification is simply not required.

Related Work. In several cases designing the “right algorithm” on graphs of bounded treewidth or pathwidth is not at all obvious. For example: Alber et al. [5] gave a $\mathcal{O}^*(4^{\text{tw}(G)})$ time algorithm for DOMINATING SET, improving over the natural $\mathcal{O}^*(9^{\text{tw}(G)})$ algorithm of Telle and Proskurowski [48]. Later, van Rooij et al. [50] observed that one could use fast subset convolution [8] to improve the running time of algorithms on graphs of bounded treewidth. Their results include a $\mathcal{O}^*(3^{\text{tw}(G)})$ algorithm for DOMINATING SET and a $\mathcal{O}^*(2^{\text{tw}(G)})$ time algorithm for PARTITION INTO TRIANGLES. Interestingly, the effect of applying subset convolution was that the running time for several graph problems on bounded treewidth graphs became the same as the running time for the problems on graphs of bounded pathwidth. However, the idea of using subset convolution in designing dynamic programming algorithm over graphs of bounded treewidth was not enough to design “optimal algorithms” for several connectivity problems such as HAMILTONIAN PATH and CONNECTED VERTEX COVER. In a seminal paper, Cygan et al. [21] introduced the method of Cut & Count and designed the first $\mathcal{O}^*(c^{\text{tw}(G)})$, where c is a fixed constant, for plethora of connectivity problems including HAMILTONIAN PATH and CONNECTED VERTEX COVER. However, the algorithm for HAMILTONIAN PATH runs in time $\mathcal{O}^*(4^{\text{tw}(G)})$, which still is the best known algorithm. Later, in a surprising result, Cygan, Kratsch, and Nederlof [20] showed that HAMILTONIAN PATH can be solved in time $\mathcal{O}^*((2 + \sqrt{2})^{\text{pw}(G)})$ on graphs of bounded pathwidth. The algorithms obtained using Cut & Count are randomized. Later, deterministic algorithms with running time $\mathcal{O}^*(c^{\text{tw}(G)})$, where c is a fixed constant, were designed for connectivity problems [10, 30].

Follow-up Work. The problems considered in this article, and the ideas used to resolve them, led to several follow-up works that showed lower bounds for concrete problems in the parameterized settings [11, 16, 17, 21]. On the other hand, Roditty and Williams [45] used SETH to show concrete lower bounds on the running time for problems solvable in polynomial time. This was followed by a long line of work in this direction [1, 2, 3, 6, 13, 51]. The work of Cygan et al. [21] that introduced the method of Cut & Count to design $\mathcal{O}^*(c^{\text{tw}(G)})$, where c is a fixed constant, for connectivity problems, also showed that the base of exponent in their algorithm are optimal unless SETH fails. Cygan, Kratsch, and Nederlof [20] showed that the running time of $\mathcal{O}^*((2 + \sqrt{2})^{\text{pw}(G)})$ for HAMILTONIAN PATH on graphs of bounded pathwidth is in fact optimal under SETH. Several other lower bounds for concrete problems were also obtained in [17]. Ideas from the current paper were recently used to design tight lower bounds for r -DOMINATING SET and CONNECTED DOMINATING SET on graphs of bounded treewidth [11]. Curticapean and Marx obtained tight lower bounds for counting perfect matchings on graphs of bounded treewidth, cliquewidth, and genus under SETH [16].

2 Preliminaries

In this section we give various definitions which we make use of in the paper. Let G be a graph with vertex set $V(G)$ and edge set $E(G)$. A graph G' is a *subgraph* of G if $V(G') \subseteq V(G)$ and $E(G') \subseteq E(G)$. For a subset $V' \subseteq V(G)$, the subgraph $G' = G[V']$ of G is called a *subgraph induced by V'* if $E(G') = \{uv \in E(G) \mid u, v \in V'\}$ and $V(G') = V'$. By $N(u)$ we denote the (open) neighborhood of u in graph G , that is, the set of all vertices adjacent to u and by $N[u] = N(u) \cup \{u\}$. Similarly, for a subset $D \subseteq V(G)$, we define $N[D] = \cup_{v \in D} N[v]$.

A *tree decomposition* of a graph G is a pair (\mathcal{X}, T) where T is a tree and $\mathcal{X} = \{X_i \mid i \in V(T)\}$ is a collection of subsets of $V(G)$ such that: **1.** $\bigcup_{i \in V(T)} X_i = V(G)$, **2.** for each edge $xy \in E(G)$, $\{x, y\} \subseteq X_i$ for some $i \in V(T)$; **3.** for each $x \in V(G)$ the set $\{i \mid x \in X_i\}$ induces a connected subtree of T . The *width* of the tree decomposition is $\max_{i \in V(T)} \{|X_i| - 1\}$. The *treewidth* of a graph G is the minimum width over all tree decompositions of G . We denote by $\text{tw}(G)$ the treewidth of graph G . If in the definition of treewidth we restrict the tree T to be a path then

we get the notion of pathwidth and denote it by $\mathbf{pw}(G)$. For our purpose we need an equivalent definition of pathwidth via *mixed search* games.

In a mixed search game, a graph G is considered as a system of tunnels. Initially, all edges are contaminated by a gas. An edge is *cleared* by placing searchers at both its end-points simultaneously or by sliding a searcher along the edge. A cleared edge is re-contaminated if there is a path from an uncleared edge to the cleared edge without any searchers on its vertices or edges. A search is a sequence of operations that can be of the following types: (a) placement of a new searcher on a vertex; (b) removal of a searcher from a vertex; (c) sliding a searcher on a vertex along an incident edge and placing the searcher on the other end. A search strategy is winning if after its termination all edges are cleared. The mixed search number of a graph G , denoted by $\mathbf{ms}(G)$, is the minimum number of searchers required for a winning strategy of mixed searching on G . Takahashi, Ueno and Kajitani [47] obtained the following relationship between $\mathbf{pw}(G)$ and $\mathbf{ms}(G)$, which we use for bounding the pathwidth of the graphs obtained in reduction.

Proposition 1 ([47]). *For a graph G , $\mathbf{pw}(G) \leq \mathbf{ms}(G) \leq \mathbf{pw}(G) + 1$.*

An instance to SAT consists of a boolean formula $\phi = C_1 \wedge \dots \wedge C_m$ over n variables $\{v_1, \dots, v_n\}$ where each clause C_i is OR of one or more literals of variables. We also denote a clause C_i by the set $\{\ell_1, \ell_2, \dots, \ell_c\}$ of its literals and denote by $|C_i|$ the number of literals in C_i . An assignment τ to the variables is an element of $\{0, 1\}^n$, and it satisfies the formula ϕ if for every clause C_i there is literal that is assigned 1 by τ . We say that a variable v_i satisfies a clause C_j if there exists a literal corresponding to v_i in $\{\ell_1, \ell_2, \dots, \ell_c\}$ and it is set to 1 by τ . A group of variables satisfy a clause C_j if there is a variable that satisfies the clause C_j . All the sections in this paper follows the following pattern: definition of the problem; statement of the lower bound; construction used in the reduction; correctness of the reduction; and the upper bound on the pathwidth of the resultant graph.

3 Independent Set

An *independent set* of a graph G is a set $S \subseteq V(G)$ such that $G[S]$ contains no edges. In the INDEPENDENT SET problem we are given a graph G and the objective is to find an independent set of maximum size.

We first sketch the main idea of the proof. We give the reduction from an arbitrary SAT instance on n variables and m clauses. The idea is to create a family of n very long paths P_1, P_2, \dots, P_n of even length, corresponding to variables x_1, x_2, \dots, x_n . Assume for now that on each of these paths the solution is allowed to make one of two choices: the independent set either contains all the odd-indexed vertices, or all the even-indexed vertices. Then for every clause we construct a clause verification gadget and attach it to some place in the family. The gadget is adjacent to paths corresponding to variables appearing in the clause, and the attachment points reflect whether the variable's appearance is positive or negative. The role of the clause gadget is to verify that the clause is satisfied. Satisfaction of the clause corresponds to the condition that at least one of the attachment points of the clause gadget needs to be *not* chosen into the constructed independent set; hence the clause gadget needs to have the following property: the behavior inside the gadget can be set optimally if and only if at least one of the attachment points is free. It is possible to construct a gadget with exactly this property, and moreover the gadget has constant pathwidth, so it does not increase much the width of the whole construction. One technical problem that we still need to overcome is the first technical assumption about the choices the solution makes on the paths P_i . It is namely not true that on a path of even length there are only two maximum-size independent sets: the odd-indexed vertices and the even-indexed vertices. The solution can first start with picking only odd-indexed vertices, then

make a gap of two vertices, and continue further with even-indexed vertices. Thus, on each path there can be one “cheat” where the solution flips from odd indices to even indices. The solution to this problem is a remarkably simple trick that is commonly used in similar reductions. We namely repeat the whole sequence of clause gadgets $n + 1$ times, which ensures that at most n copies are spoiled by possible cheats, and hence at least one of the copies is attached to area where no cheat happens, and hence the behavior of the solution on the paths P_i correctly encodes some satisfying assignment of the variable set. This concludes the sketch and we move towards giving the formal proof.

Theorem 1. *If INDEPENDENT SET can be solved in $\mathcal{O}^*((2 - \epsilon)^{\text{tw}(G)})$ for some $\epsilon > 0$ then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Construction. Given an instance ϕ of SAT, we construct a graph G as follows (see Figure 1). We assume that every clause has an even number of variables: if not, we can add a single variable to all odd size clauses and force this variable to false. First we describe the construction of clause gadgets. For a clause $C = \{\ell_1, \ell_2, \dots, \ell_c\}$, we introduce a gadget \widehat{C} as follows. We take two paths, $CP = cp_1, cp_2 \dots, cp_c$ and $CP' = cp'_1, cp'_2 \dots cp'_c$ having c vertices each, and connect cp_i with cp'_i for every i . For each literal ℓ_i , we introduce a vertex ℓ_i in \widehat{C} and make it adjacent to cp_i and cp'_i . Finally we add two vertices c_{start} and c_{end} , such that c_{start} is adjacent to cp_1 and c_{end} is adjacent to cp_c . Observe that the size of the maximum independent set of \widehat{C} is $c + 2$. Also, since c is even, any independent set of size $c + 2$ in \widehat{C} must contain at least one vertex in $C = \{\ell_1, \ell_2, \dots, \ell_c\}$. Finally, notice that for any i , there is an independent set of size $c + 2$ in \widehat{C} that contains ℓ_i and none of ℓ_j for $j \neq i$.

We first construct a graph G_1 . We introduce n paths P_1, \dots, P_n , each path has $2m$ vertices. Let the vertices of the path P_i be $p_i^1 \dots p_i^{2m}$. The path P_i corresponds to the variable v_i . For every clause C_j of ϕ , we introduce a gadget \widehat{C}_j . Now, for every variable v_i , if v_i occurs positively in C_j , we add an edge between p_i^{2j} and the literal corresponding to v_i in \widehat{C}_j . If v_i occurs negatively in C_j , we add an edge between p_i^{2j-1} and the literal corresponding to v_i in \widehat{C}_j . Now we construct the graph G as follows. We take $n + 1$ copies of G_1 , call them G_1, \dots, G_{n+1} . For every $i \leq n$, we connect G_i and G_{i+1} by connecting p_j^{2m} in G_i with p_j^1 in G_{i+1} for every $j \leq n$. This way, the paths P_j in each of the n copies G_i together form a long path of $2m(n + 1)$ vertices. This concludes the construction of G .

Lemma 1. *If ϕ is satisfiable, then G has an independent set of size $(mn + \sum_{i \leq m} (|C_i| + 2))(n + 1)$.*

Proof. Consider a satisfying assignment to ϕ . We construct an independent set I in G . For every variable v_i , if v_i is set to true, then pick all the vertices on odd positions from all copies of P_i , that is p_i^1, p_i^3, p_i^5 and so on. If v_i is false then pick all the vertices on even positions from all copies of P_i , that is p_i^2, p_i^4, p_i^6 and so on. It is easy to see that this is an independent set of size $mn(n + 1)$ containing vertices from all the paths. We will now consider the gadget \widehat{C}_j corresponding to a clause C_j . We will only consider the copy of \widehat{C}_j in G_1 as the other copies can be dealt identically. Let us choose a true literal ℓ_a in C_j and let v_i be the corresponding variable. Consider the vertex ℓ_a in \widehat{C}_j . If v_i occurs positively in C_j , then v_i is true. Then I does not contain p_i^{2j} , the only neighbour of ℓ_a outside of \widehat{C}_j . On the other hand if v_i occurs negatively in C_j , then v_i is false. In this case I does not contain p_i^{2j-1} , the only neighbour of ℓ_a outside of \widehat{C}_j . There is an independent set of size $|C_j| + 2$ in \widehat{C}_j that contains ℓ_a and none out of ℓ_b for any $b \neq a$. We add this independent set to I and proceed in this manner for every clause gadget. By the end of the process $(\sum_{i \leq m} (|C_i| + 2))(n + 1)$ vertices from clause gadgets are added to I , yielding that the size of I is $(mn + \sum_{i \leq m} (|C_i| + 2))(n + 1)$, concluding the proof. \square

Lemma 2. *If G has an independent set of size $(mn + \sum_{i \leq m} (|C_i| + 2))(n + 1)$, then ϕ is satisfiable.*




Figure 1: Reduction to INDEPENDENT SET: clause gadget \widehat{C}_j attached to the n paths representing the variables.

Proof. Consider an independent set of G of size $(mn + \sum_{i \leq m} |C_i| + 2)(n + 1)$. Set I can contain at most m vertices from each copy of P_i for every $i \leq n$ and at most $|C_j| + 2$ vertices from each copy of the gadget C_j . Since I must contain at least that many vertices from each path and clause gadget in order to contain at least $(mn + \sum_{i \leq m} |C_i| + 2)(n + 1)$ vertices, it follows that I has exactly m vertices in each copy of each path P_i and exactly $|C_j| + 2$ vertices in each copy of each clause gadget \widehat{C}_j . For a fixed j , consider the $n + 1$ copies of the path P_j . Since P_j in G_i is attached to P_j in G_{i+1} , these $n + 1$ copies of P_j together form a path P having $2m(n + 1)$ vertices. Since $|I \cap P| = m(n + 1)$ it follows that $I \cap P$ must contain every second vertex of P , except possibly in one position where $I \cap P$ skips two vertices of P . There are only n paths and $n + 1$ copies of G_1 , hence the pigeon-hole principle implies that in some copy G_y of G_1 , I contains every second vertex on every path P_i . From now onwards we only consider such a copy G_y .

In G_y , for every $i \leq n$, I contains every second vertex of P_i . We make an assignment to the variables of ϕ as follows. If I contains all the odd numbered vertices of P_i then v_i is set to true, otherwise I contains all the even numbered vertices of P_i and v_i is set to false. We argue that this assignment satisfies ϕ . Indeed, consider any clause C_j , and look at the gadget \widehat{C}_j . We know that I contains $|C_j| + 2$ vertices from \widehat{C}_j and hence I must contain a vertex ℓ_a in \widehat{C}_j corresponding to a literal of C_j . Suppose ℓ_a is a literal of v_i . Since I contains ℓ_a , if ℓ_a occurs positively in C_j , then I can not contain p_i^{2j} and hence v_i is true. Similarly, if ℓ_a occurs negatively in C_j then I can not contain p_i^{2j-1} and hence v_i is false. In both cases v_i satisfies C_j and hence all clauses of ϕ are satisfied by the assignment. \square

Lemma 3. $\text{pw}(G) \leq n + 4$.

Proof. We give a mixed search strategy to clean G using $n + 3$ searchers. For every i we place a searcher on the first vertex of P_i in G_1 . The n searchers slide along the paths P_1, \dots, P_n in m rounds. In round j each searcher i starts on p_i^{2j-1} . Then, for every variable v_i that occurs positively in C_j , the searcher i slides forward to p_i^{2j} . Observe that at this point there is a searcher on every neighbour of the gadget \widehat{C}_j . This gadget can now be cleaned with 3 additional searchers. After \widehat{C}_j is clean, the additional 3 searchers are removed, and each of the n searchers on the paths P_1, \dots, P_n slides forward along these paths, such that searcher i stands on $p_i^{2(j+1)}$. At that point, the next round commences. When the searchers have cleaned G_1 they slide onto the first vertex of $P_1 \dots P_n$ in G_2 . Then they proceed to clean G_2, \dots, G_{n+1} in the same way that G_1 was cleaned. Now applying Proposition 1 we get that $\text{pw}(G) \leq n + 4$. \square

The construction, together with Lemmata 1, 2 and 3 proves Theorem 1.

4 Dominating Set

A *dominating set* of a graph G is a set $S \subseteq V(G)$ such that $V(G) = N[S]$. In the DOMINATING SET problem we are given a graph G and the objective is to find a dominating set of minimum size.

The basic idea for this reduction is similar to the one for INDEPENDENT SET. However, we need one more new idea here, which will also be used in other reductions. We group variables into an appropriate number of groups of size at most $\beta = \lfloor \log 3^p \rfloor$, where p is a constant depending only on ϵ . Then, for every group we make a gadget such that an assignment on the group should correspond to a selection on the gadget. These group gadgets are then connected to clause gadgets so that every assignment on the group that satisfies the clause results in some desired outcome.




Figure 2: Reduction to DOMINATING SET: group gadget \widehat{B} . The set S is shown by the circled vertices.

Theorem 2. *If DOMINATING SET can be solved in $\mathcal{O}^*((3 - \epsilon)^{\text{pw}(G)})$ time for some $\epsilon > 0$ then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Construction. Given $\epsilon < 1$ and an instance ϕ to SAT we construct a graph G as follows. We first choose an integer p depending only on ϵ . Exactly how p is chosen will be discussed in the proof of Theorem 2. We group the variables of ϕ into groups F_1, F_2, \dots, F_t , each of size at most $\beta = \lfloor \log 3^p \rfloor$. Hence $t = \lceil n/\beta \rceil$. We now proceed to describe a “group gadget” \widehat{B} , which is central in our construction.

To build the group gadget \widehat{B} , we introduce p paths P_1, \dots, P_p , where the path P_i contains the vertices p_i^1, p_i^2 and p_i^3 (see Figure 2). To each path P_i we attach two guards g_i and g'_i , both of which are neighbours to p_i^1, p_i^2 and p_i^3 . When the gadgets are attached to each other, the guards will not have any neighbours outside of their own gadget \widehat{B} , and will ensure that at least one vertex out of p_i^1, p_i^2 and p_i^3 are chosen in any minimum size dominating set of G . Let P be the vertex set containing all the vertices on the paths P_1, \dots, P_p . For every subset S of P that picks *exactly one* vertex from each path P_i , we introduce two vertices x_S and x'_S , where x_S is adjacent to all vertices of $P \setminus S$ (all those vertices that are on paths and not in S) and x'_S is only adjacent to x_S . We conclude the construction of \widehat{B} by making all the vertices x'_S (for every set S) adjacent to each other, that is making them into a clique, and adding a guard x adjacent to x'_S for every set S . In other words, the x'_S 's together with x form a clique and all the neighbors of x reside in this clique.

We construct the graph G as follows (see Figure 3). For every group F_i of variables, we introduce $m(2pt + 1)$ copies of the gadget \widehat{B} , call them \widehat{B}_i^j for $1 \leq j \leq m(2pt + 1)$. We can imagine these $t \cdot m(2pt + 1)$ gadgets arranged in t rows and $m(2pt + 1)$ columns, with the columns being divided into $2pt + 1$ regions of m columns each. For every fixed $i \leq t$, we connect the




Figure 3: Reduction to DOMINATING SET: arranging the group gadgets. Note that $x = ml + j$, thus \hat{c}_j^ℓ is attached to vertices in $\hat{B}_1^x, \dots, \hat{B}_t^x$.

gadgets $\hat{B}_i^1, \hat{B}_i^2, \dots, \hat{B}_i^{m(2pt+1)}$ in a path-like manner. In particular, for every $j < m(2pt+1)$ and every $\ell \leq p$ we make an edge between p_ℓ^3 in the gadget \hat{B}_i^j with p_ℓ^1 in the gadget \hat{B}_i^{j+1} . Now we introduce two new vertices h and h' , with h adjacent to h' , p_j^1 in \hat{B}_i^1 for every $i \leq t, j \leq p$ and to p_j^3 in $\hat{B}_i^{m(2pt+1)}$ for every $i \leq t, j \leq p$. That is, for all $1 \leq i \leq t$, h is adjacent to the first and last vertices of “long paths” obtained after connecting the gadgets $\hat{B}_i^1, \hat{B}_i^2, \dots, \hat{B}_i^{m(2pt+1)}$ in a path-like manner.

For every $1 \leq i \leq t$ and to every assignment of the variables in the group F_i , we designate a subset S of P in the gadget \hat{B} that picks exactly one vertex from each path P_j . Since there are at most 2^β different assignments to the variables in F_i , and there are $3^p \geq 2^\beta$ such sets S , we can assign a *unique* set to each assignment. Of course, the same set S can correspond to one assignment of the group F_1 and some other assignment of the group F_2 . Recall that the clauses of ϕ are C_1, \dots, C_m . For every clause C_j we introduce $2pt+1$ vertices \hat{c}_j^ℓ , one for each $0 \leq \ell < 2pt+1$, corresponding to the $2pt+1$ regions. The vertex \hat{c}_j^ℓ will be connected to the gadgets $\hat{B}_i^{m\ell+j}$ for every $1 \leq i \leq t$ (which appear in the ℓ -th region). In particular, for every assignment of the variables in the group F_i that satisfy the clause C_j , we consider the

subset S of P that corresponds to the assignment. For every $0 \leq \ell < 2pt + 1$, we make x'_S in $\widehat{B}_i^{m\ell+j}$ adjacent to \widehat{c}_j^ℓ . The best way to view this is that every clause C_j has $2pt + 1$ private gadgets in the i -row, $\widehat{B}_i^j, \widehat{B}_i^{m+j}, \dots, \widehat{B}_i^{m2pt+j}$, one in each region. Now we have $2pt + 1$ vertices corresponding to the clause C_j , each connected to one of these gadgets. This concludes the construction of G .

Lemma 4. *If ϕ has a satisfying assignment, then G has a dominating set of size $(p+1)tm(2pt+1) + 1$.*

Proof. Given a satisfying assignment to ϕ , we construct a dominating set D of G that contains the vertex h and *exactly* $p + 1$ vertices in each gadget \widehat{B}_i^j . For each group F_i of variables we consider the set S that corresponds to the restriction of the assignment to the variables in F_i . From each gadget \widehat{B}_i^j we add the set S to D and also the vertex x'_S to D . It remains to argue that D is indeed a dominating set. Clearly the size is bounded by $(p + 1)tm(2pt + 1) + 1$, as the number of gadgets is $tm(2pt + 1)$.

For a fixed $i \leq t$ and j consider the vertices on the path P_j in the gadgets \widehat{B}_i^ℓ for every $\ell \leq m(2pt + 1)$. Together these vertices form a path of length $3m(2pt + 1)$ and every third vertex of this path is in S . Thus, all vertices on this path are dominated by other vertices on the path, except perhaps for the first and last one. Both these vertices, however, are dominated by h .

Now, fix some $i \leq t$ and $\ell \leq m(2pt + 1)$ and consider the gadget \widehat{B}_i^ℓ . Since D contains some vertex on the path P_j , we have that for every j both g_j and g'_j are dominated. Furthermore, for every set S^* not equal to S that picks exactly one vertex from each P_j , vertex x_{S^*} is dominated by some vertex on some P_j —namely by all vertices in $S \setminus S^* \neq \emptyset$. The last assertion follows since x_{S^*} is connected to all the vertices on the paths except S^* . On the other hand, x_S is dominated by x'_S , and x'_S also dominates all the other vertices x'_{S^*} for $S^* \neq S$, as well as the guard x .

The only vertices not yet accounted for are the vertices \widehat{c}_j^ℓ for every $j \leq m$ and $\ell < 2pt + 1$. Fix a j and a ℓ and consider the clause C_j . This clause contains a literal set to true, and this literal corresponds to a variable in the group F_i for some $i \leq t$. Of course, the assignment to F_i satisfies C_j . Let S be the set corresponding to this assignment of F_i . By the construction of D , the dominating set contains x'_S in $\widehat{B}_i^{m\ell+j}$ and x'_S is adjacent to \widehat{c}_j^ℓ . This concludes the proof. \square

Lemma 5. *If G has a dominating set of size $(p + 1)tm(2pt + 1) + 1$, then ϕ has a satisfying assignment.*

Proof. Let D be a dominating set of G of size at most $(p + 1)tm(2pt + 1) + 1$. Since D must dominate h' , without loss of generality we can assume that D contains h . Furthermore, inside every gadget \widehat{B}_i^ℓ , D must dominate all the guards, namely g_j and g'_j for every $j \leq p$, and also x . Thus D contains at least $p + 1$ vertices from each gadget \widehat{B}_i^ℓ which in turn implies that D contains exactly $p + 1$ vertices from each gadget \widehat{B}_i^ℓ . The only way D can dominate g_j and g'_j for every j and in addition dominate x with only $p + 1$ vertices is if D has one vertex from each P_j , $j \leq p$ and in addition contains some vertex in $N[x]$. Let S be $D \cap P$ in \widehat{B}_i^ℓ . Observe that x_S is not dominated by $D \cap S$. The only vertex in $N[x]$ that dominates x_S is x'_S and hence D contains x'_S .

Now we want to show that for every $1 \leq i \leq t$ there exists *one* $0 \leq \ell \leq 2tp$ such that for fixed i , $D \cap P$ is same in all the gadgets $\widehat{B}_i^{m\ell+r}$ for every $1 \leq r \leq m$, i.e., it is the same in every gadget of the i -th row in the ℓ -th region. Consider a gadget \widehat{B}_i^ℓ and its follower, $\widehat{B}_i^{\ell+1}$. Let S be $D \cap P$ in \widehat{B}_i^ℓ and S' be $D \cap P$ in $\widehat{B}_i^{\ell+1}$. Observe that if S contains p_j^a in \widehat{B}_i^ℓ and p_j^b in $\widehat{B}_i^{\ell+1}$ then we must have $b \leq a$. We call a consecutive pair *bad* if for some $j \leq p$, D contains

p_j^a in \widehat{B}_i^ℓ and p_j^b in $\widehat{B}_i^{\ell+1}$ and $b < a$. Hence for a fixed i , we can at most have $2p$ consecutive bad pairs, spoiling at most $2p$ regions. Now we mark all the bad pairs that occur among the gadgets corresponding to some F_i . This way we can mark only $2tp$ bad pairs. Thus, by the pigeon hole principle, there exists an $\ell \in \{0, \dots, 2tp\}$ such that there are no bad pairs in $\widehat{B}_i^{m\ell+r}$ for all $1 \leq i \leq t$ and $1 \leq r \leq m$.

We make an assignment ϕ by reading off $D \cap P$ in each gadget $\widehat{B}_i^{m\ell+1}$. In particular, for every group F_i , we consider $S = D \cap P$ in the gadget $\widehat{B}_i^{m\ell+1}$. This set S corresponds to an assignment of F_i , and this is the assignment of F_i that we use. It remains to argue that every clause C_r is satisfied by this assignment.

Consider the vertex \widehat{c}_ℓ^r . We know that it is dominated by some x'_S in a gadget $\widehat{B}_i^{m\ell+r}$. The set S corresponds to an assignment of F_i that satisfies the clause C_r . Because $D \cap P$ remains unchanged in all gadgets from $\widehat{B}_i^{m\ell+1}$ to $\widehat{B}_i^{m\ell+r}$, this is exactly the assignment ϕ restricted to the group F_i . This concludes the proof. \square

Lemma 6. $\text{pw}(G) \leq tp + \mathcal{O}(3^p)$

Proof. We give a mixed search strategy to clean the graph with $tp + \mathcal{O}(3^p)$ searchers. For a gadget \widehat{B} we call the vertices p_j^1 and p_j^3 , $1 \leq j \leq p$, as *entry vertices* and *exit vertices* respectively. We search the graph in $m(2tp + 1)$ rounds. In the beginning of round ℓ there are searchers on the entry vertices of the gadgets \widehat{B}_i^ℓ for every $i \leq t$. Let $1 \leq a \leq m$ and $0 \leq b < 2tp + 1$ be integers such that $\ell = a + mb$. We place a searcher on \widehat{c}_a^b . Then, for each i between 1 and p in turn we first put searchers on all vertices of \widehat{B}_i^ℓ and then remove all the searchers from \widehat{B}_i^ℓ except for the ones standing on the exit vertices. After all gadgets $\widehat{B}_1^\ell \dots \widehat{B}_t^\ell$ have been cleaned in this manner, we can remove the searcher from \widehat{c}_a^b . To commence the next round, the searchers slide from the exit positions of \widehat{B}_i^ℓ to the entry positions of $\widehat{B}_i^{\ell+1}$ for every i . In total, at most $tp + |V(\widehat{B})| + 1 \leq tp + \mathcal{O}(3^p)$ searchers are used simultaneously. This together with Proposition 1 give the desired upperbound on the pathwidth. \square

Proof (of Theorem 2). Suppose DOMINATING SET can be solved in $\mathcal{O}^*((3-\epsilon)^{\text{pw}(G)}) = \mathcal{O}^*(3^{\lambda \text{pw}(G)})$ time, where $\lambda = \log_3(3-\epsilon) < 1$. We choose p large enough such that $\lambda \cdot \frac{p}{\lfloor p \log 3 \rfloor} = \frac{\delta'}{\log 3}$ for some $\delta' < 1$. Given an instance of SAT, we construct an instance of DOMINATING SET using the above construction and the chosen value of p . Then we solve the DOMINATING SET instance using the $\mathcal{O}^*(3^{\lambda \text{pw}(G)})$ time algorithm. Correctness is ensured by Lemmata 4 and 5. Lemma 6 yields that the total time taken is upper bounded by $\mathcal{O}^*(3^{\lambda \text{pw}(G)}) \leq \mathcal{O}^*(3^{\lambda(tp+f(\lambda))}) \leq \mathcal{O}^*(3^{\lambda \frac{np}{\lfloor p \log 3 \rfloor}}) \leq \mathcal{O}^*(3^{\delta' \frac{n}{\log 3}}) \leq \mathcal{O}^*(2^{\delta'' n}) = \mathcal{O}^*((2-\delta)^n)$, for some $\delta'', \delta < 1$. This concludes the proof. \square

5 Max Cut

A *cut* in a graph G is a partition of $V(G)$ into V_0 and V_1 . The *cut-set* of the cut is the set of edges whose one end point is in V_0 and the other in V_1 . We say that an edge is *crossing* this cut if it has one endpoint in V_0 and one in V_1 , that is, the edge is in the cut-set. The *size* of the cut is the number of edges in G which are crossing this cut. If the edges of G have positive integer weights, then the *weight* of the cut is the sum of the weights of edges that are crossing the cut. In the MAX CUT problem, we are given a graph G together with an integer t and asked whether there is a cut of G of size at least t . In the WEIGHTED MAX CUT problem every edge has a positive integer weight and the objective is to find a cut of weight at least t .

Theorem 3. *If MAX CUT can be solved in $\mathcal{O}^*((2-\epsilon)^{\text{pw}(G)})$ for some $\epsilon > 0$, then SAT can be solved in $\mathcal{O}^*((2-\delta)^n)$ time for some $\delta > 0$.*




Figure 4: Reduction to MAX CUT. The dashed edges have weight 1 and all the other edges have weight $3n$. The odd and even positions of the paths P_j are shown by black and white, respectively. As an example, we show potential connections corresponding to the clauses $C_1 = (v_1 \vee \bar{v}_2 \vee v_4)$ and $C_2 = (\bar{v}_1 \vee v_3 \vee \bar{v}_5)$.

Construction. Given an instance ϕ of SAT, we first construct an instance G_w of WEIGHTED MAX CUT as follows. We later explain how to obtain an instance of unweighted MAX CUT from here.

We start with introducing a vertex x_0 . Without loss of generality, we will assume that $x_0 \in V_0$ in every solution. We introduce a vertex \hat{v}_i for each variable v_i . For every clause C_j , we create a gadget as follows. We introduce a path \hat{P}_j having $4|C_j|$ vertices. All the edges on \hat{P}_j have weight $3n$. Now, we make the first and last vertex of \hat{P}_j adjacent to x_0 with an edge of weight $3n$. Thus the path \hat{P}_j plus the edges from the first and last vertex of \hat{P}_j to x_0 form an odd cycle \hat{C}_j . We will say that the first, third, fifth, etc, vertices are on *odd positions* on \hat{P}_j while the remaining vertices are on *even positions*. For every variable v_i that appears positively in C_j , we select a vertex p at an even position (but not the last vertex) on \hat{P}_j and make \hat{v}_i adjacent to p and p 's successor on \hat{P}_j with edges of weight 1. For every variable v_i that appears negatively in C_j we select a vertex p at an odd position on \hat{P}_j and make \hat{v}_i adjacent to p and p 's successor on \hat{P}_j with edges of weight 1. We make sure that each vertex on \hat{P}_j receives an edge at most once in this process. There are more than enough vertices on \hat{P}_j to accommodate all the edges incident to vertices corresponding to variables in the clause C_j . We create such a gadget for each clause and set $t = m + (12n + 1) \sum_{j=1}^m |C_j|$. This concludes the construction.

Lemma 7. *If ϕ is satisfiable, then G_w has a cut of weight at least t .*

Proof. Suppose ϕ is satisfiable. We put x_0 in V_0 and for every variable v_i we put \hat{v}_i in V_1 if v_i is true and \hat{v}_i in V_0 if v_i is false. For every clause C_j we proceed as follows. Let us choose a true literal of C_j and suppose that this literal corresponds to a vertex p_j on \hat{P}_j . We put the first vertex on \hat{P}_j in V_1 , the second in V_0 and then we proceed along \hat{P}_j putting every second vertex into V_1 and V_0 until we reach p_j . The successor p'_j of p_j on \hat{P}_j is put into the same set as p_j .

Then we continue along \widehat{P}_j putting every second vertex in V_1 and V_0 . Notice that even though C_j may contain more than one literal that is set to true, we only select one vertex p_j from the path \widehat{P}_j and put p_j and its successor on the same side of the partition. It remains to argue that this cut has weight at least t .

For every clause C_j all edges on the path \widehat{P}_j except for $p_j p'_j$ are crossing, and the two edges to x_0 from the first and last vertex of \widehat{P}_j are crossing as well. These edges contribute $12n|C_j|$ to the weight of the cut. We know that p_j corresponds to a literal that is set to true, and this literal corresponds to a variable v_i . If v_i occurs positively in C_j , then $\widehat{v}_i \in V_1$ and p_j is on an even position of \widehat{P}_j . Thus both p_j and its successor p'_j are in V_0 and hence both $\widehat{v}_i p_j$ and $\widehat{v}_i p'_j$ are crossing, contributing 2 to the weight of the cut. For each of the remaining variables $v_{i'}$ appearing in C_j , one of the two neighbours of $\widehat{v}_{i'}$ on \widehat{P}_j appear in V_0 and one in V_1 , so exactly one edge from $v_{i'}$ to \widehat{P}_j is crossing. Thus the total weight of the cut is $t = \sum_{j=1}^m (12n|C_j| + |C_j| + 1) = m + (12n + 1) \sum_{j=1}^m |C_j|$. This completes the proof. \square

Lemma 8. *If G_w has a cut of weight at least t , then ϕ is satisfiable.*

Proof. Let (V_0, V_1) be a cut of G of maximum weight, hence the weight of this cut is at least t . Without loss of generality, let $x_0 \in V_0$. For every clause C_j , at least one edge of the odd cycle \widehat{C}_j is not crossing. If more than one edge of this cycle is not crossing, then the total weight of the cut edges incident to the path \widehat{P}_j is at most $3n(4|C_j| - 1) + 2n < 12n|C_j|$. In this case, we could change the partition (V_0, V_1) such that all edges of \widehat{P}_j are crossing and the first vertex of \widehat{P}_j is in V_1 . Using the new partition the weight of the crossing edges in the cycle \widehat{C}_j is at least $12n|C_j|$ and the edges not incident to \widehat{P}_j are unaffected by the changes. This contradicts that (V_0, V_1) was a maximum weight cut. Thus it follows that exactly one edge of \widehat{C}_j is not crossing.

Given the cut (V_0, V_1) , we set each variable v_i to true if $\widehat{v}_i \in V_1$ and v_i to false otherwise. Consider a clause C_j and a variable v_i that appears in C_j . Let uv be the edge of \widehat{C}_j that is not crossing. If there is a vertex \widehat{v}_i adjacent to both u and v , then it is possible that both $\widehat{v}_i u$ and $\widehat{v}_i v$ are crossing. For every other variable $v_{i'}$ in C_j , at most one of the edges from $\widehat{v}_{i'}$ to \widehat{P}_j is crossing. Thus, the weight of the edges that are crossing in the gadget \widehat{C}_j is at most $(12n + 1)|C_j| + 1$. Hence, to find a cut-set of weight at least t in G , we need to have crossing edges in \widehat{C}_j with sum of their weights exactly equal to $12n|C_j| + |C_j| + 1$. It follows that there is a vertex \widehat{v}_i adjacent to both u and v such that both $\widehat{v}_i u$ and $\widehat{v}_i v$ are crossing.

If v_i occurs in C_j positively, then u is on an even position and hence, $u \in V_0$. Since $\widehat{v}_i u$ is crossing it follows that v_i is true and C_j is satisfied. On the other hand, if v_i occurs in C_j negated then u is on an odd position and hence, $u \in V_1$. Since $\widehat{v}_i u$ is crossing it follows that v_i is false and C_j is satisfied. As this holds for each clause individually, this concludes the proof. \square

For every edge $e \in E(G_w)$, let w_e be the weight of e in G_w . We construct an unweighted graph G from G_w by replacing every edge $e = uv$ by w_e paths from u to v on three edges. Let W be the sum of the edge weights of all edges in G_w .

Lemma 9. *G has a cut of size $2W + t$ if and only if G_w has a cut of weight at least t .*

Proof. Given a partition of $V(G_w)$, we partition $V(G)$ as follows. The vertices of G that also are vertices of $V(G_w)$ are partitioned in the same way as in $V(G_w)$. On each path of length 3, if the endpoints of the path are in different sets we can partition the middle vertices of the path such that all edges are cut. If the endpoints are in the same set we can only partition the middle vertices such that 2 out of the 3 edges are cut. The reverse direction is similar. \square

Lemma 10. $\text{pw}(G) \leq n + 5$.

Proof. We give a search strategy to clean G with $n + 5$ searchers. We place one searcher on each vertex \widehat{v}_i and one searcher on x_0 . Then one can search the gadgets \widehat{C}_j one by one. In G_w it is sufficient to use 2 searchers for each \widehat{C}_j , whereas in G after the edges have been replaced by multiple paths on three edges, we need 4 searchers. This combined with Proposition 1 gives the desired upper bound on the pathwidth of the graph. \square

The construction, together with Lemmata 7, 8, 9 and 10 proves Theorem 3.

6 Graph Coloring

A q -coloring of G is a function $\mu : V(G) \rightarrow [q]$. A q -coloring μ of G is *proper* if for every edge $uv \in E(G)$ we have $\mu(u) \neq \mu(v)$. In the q -COLORING problem we are given as input a graph G and the objective is to decide whether G has a proper q -coloring. In the LIST COLORING problem, every vertex v is given a list $L(v) \subseteq [q]$ of admissible colors. A *proper list coloring* of G is a function $\mu : V(G) \rightarrow [q]$ such that μ is a proper coloring of G that satisfies $\mu(v) \in L(v)$ for every $v \in V(G)$. In the q -LIST COLORING problem we are given a graph G together with a list $L(v) \subseteq [q]$ for every vertex v . The task is to determine whether there exists a proper list coloring of G .

A *feedback vertex set* of a graph G is a set $S \subseteq V(G)$ such that $G \setminus S$ is a forest; we denote by $\mathbf{fvs}(G)$ the size of the smallest such set. It is well-known that $\mathbf{tw}(G) \leq \mathbf{fvs}(G) + 1$. Unlike the other sections, where we give lower bounds for algorithms parameterized by $\mathbf{pw}(G)$, the following theorem gives also a lower bound for algorithms parameterized by $\mathbf{fvs}(G)$. Such a lower bound follows very naturally from the construction we are doing here, but not from the constructions in the other sections. It would be interesting to explore whether it is possible to prove tight bounds parameterized by $\mathbf{fvs}(G)$ for the problems considered in the other sections.

Theorem 4. *Let q be a fixed positive integer. If q -COLORING can be solved in $\mathcal{O}^*((q - \epsilon)^{\mathbf{fvs}(G)})$ or $\mathcal{O}^*((q - \epsilon)^{\mathbf{pw}(G)})$ time for some $\epsilon > 0$, then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Construction. We will show the result for LIST COLORING first, and then give a simple reduction that demonstrates that q -COLORING can be solved in $\mathcal{O}^*((q - \epsilon)^{\mathbf{fvs}(G)})$ time if and only if q -LIST COLORING can.

Depending on ϵ and q we choose a parameter p . Now, given an instance ϕ to SAT we will construct a graph G with a list $L(v)$ for every v , such that G has a proper list-coloring if and only if ϕ is satisfiable. Throughout the construction we will call color 1-*red*, color 2-*white* and color 3-*black*.

We start by grouping the variables of ϕ into t groups F_1, \dots, F_t of size at most $\lceil \log q^p \rceil$. Thus $t = \lceil \frac{n}{\lceil \log q^p \rceil} \rceil$. We will call an assignment of truth values to the variables in a group F_i a *group assignment*. We will say that a group assignment satisfies a clause C_j of ϕ if C_j contains at least one literal which is set to true by the group assignment. Notice that C_j can be satisfied by a group assignment of a group F_i , even though C_j also contains variables that are not in F_i .

For each group F_i , we introduce a set V_i of p vertices v_i^1, \dots, v_i^p . The vertices in V_i get full lists, that is, they can be colored by any color in $[q]$. The coloring of the vertices in V_i will encode the group assignment of F_i . There are $q^p \geq 2^{|F_i|}$ possible colorings of V_i . Thus, to each possible group assignment of F_i we attach a unique coloring of V_i . Notice that some colorings of V_i may not correspond to any group assignments of F_i .

For each clause C_j of ϕ , we introduce a gadget \widehat{C}_j . The main part of \widehat{C}_j is a long path \widehat{P}_j that has one vertex for each group assignment that satisfies C_j . Notice that there are at most tq^p possible group assignments, and that q and p are constants independent of the input ϕ . The list of every vertex on \widehat{P}_j is $\{\text{red, white, black}\}$. We attach two vertices p_j^{start} and p_j^{end} to the




Figure 5: Reduction to q -COLORING: the way the connector connects a vertex v_i^ℓ with v for a particular “bad color” $x \in [q] \setminus \{\mu_i(v_i^\ell)\}$. The left side shows the case $x = \text{red} = 1$, the right side $x = 2$ ($q = 4$).

start and end of \widehat{P}_j respectively, and the two vertices are not counted as vertices of the path \widehat{P}_j itself. The list of p_j^{start} is $\{\text{white}\}$. If $|V(\widehat{P}_j)|$ is even, then the list of p_j^{end} is $\{\text{white}\}$, whereas if $|V(\widehat{P}_j)|$ is odd then the list of p_j^{end} is $\{\text{black}\}$. The intention is that to properly color \widehat{P}_j , one needs to use the color red at least once, and that once is sufficient. The position of the red-colored vertex on the path \widehat{P}_j encodes how the clause C_j is satisfied.

For every vertex v on \widehat{P}_j , we proceed as follows. The vertex v corresponds to some group assignment to F_i that satisfies the clause C_j . This assignment in turn corresponds to a coloring of the vertices of V_i . Let this coloring be μ_i . We build a *connector* whose role is to enforce that v can be red only if coloring μ_i appears on V_i . To build the connector, for each vertex $v_i^\ell \in V_i$ and color $x \in [q] \setminus \{\mu_i(v_i^\ell)\}$ we do the following to enforce that if v is red, then v_i^ℓ *cannot* have color x (see Figure 5).

- If x is red, then we introduce one vertex w_y for every color y except for red. We make w_y adjacent to v_i^ℓ and the list of w_y is $\{\text{red}, y\}$. Then we introduce a vertex w that is adjacent to v and to all vertices w_y . The list of w is all of $[q]$.
- If x is not red, we introduce two vertices w_y and w'_y for each color y except for red. We make w_y adjacent to v_i^ℓ and w'_y adjacent to w_y . The list of w_y is $\{x, \text{red}\}$ while the list of w'_y is $\{y, \text{red}\}$. Finally, we introduce a vertex w adjacent to v and to w'_y for all y . The list of w is all of $[q]$.

Notice that in the above construction we have reused the names w , w_y and w'_y for many different vertices: in each connector, there is a separate vertex w for each vertex $v_i^\ell \in V_i$ and color $x \in [q] \setminus \{\mu_i(v_i^\ell)\}$. Building a connector for each vertex v on \widehat{P}_j concludes the construction of the clause gadget \widehat{C}_j , and creating one such gadget for each clause concludes the construction of G (see Figure 6). The following lemma, summarizes the most important properties of the connector:

Lemma 11. *Consider the connector corresponding to a vertex v on \widehat{P}_j and a coloring μ_i of V_i .*

1. *Any coloring on V_i and any color $c \in \{\text{white}, \text{black}\}$ on v can be extended to the rest of the connector.*




Figure 6: Reduction to q -COLORING. The t groups of vertices V_1, \dots, V_t represent the t groups of variables F_1, \dots, F_t (each of size $\lceil \log q^p \rceil$). Each vertex of the clause path \widehat{P}_j is connected to one group V_i via a connector (multiple vertices on the path can be connected to the same group).

2. Coloring μ_i on V_i and any color $c \in \{\text{red, white, black}\}$ on v can be extended to the rest of the connector.
3. In any coloring of the connector, if v is red, then μ_i appears on V_i .

Proof. 1. For each vertex $v_i^\ell \in V_i$ and color $x \in [q] \setminus \{\mu_i(v_i^\ell)\}$ we do the following.

- If x is red, then in the construction of \widehat{C}_j we introduced a vertex w_y with list $\{y, \text{red}\}$ for every color $y \neq \text{red}$ adjacent to v_i^ℓ , and a vertex w with list $[q]$ adjacent to w_y for every $y \neq \text{red}$. If v_i^ℓ is colored red, then we color each vertex w_y with y and w with red. Notice that w is adjacent to v , but v is colored either white or black, so it is safe to color w red. If, on the other hand, v_i^ℓ is not colored red, we can color w_y red for every y . Then all the neighbours of w have been colored with red, except for v which has been colored white or black. Thus it is safe to color w with the color out of black and white which was not used to color v .
- If x is not red, then in the construction of \widehat{C}_j we introduced two vertices w_y and w'_y for each color y except for red, and also introduced a vertex w . The vertices w_y are adjacent to v_i^ℓ and for every $y \neq \text{red}$, the vertex w'_y is adjacent to w_y . Finally, w is adjacent to all the vertices w'_y and to v . For every y the list of w_y is $\{x, \text{red}\}$ while the list of w'_y is $\{y, \text{red}\}$. The list of w is $[q]$. If v_i^ℓ is colored with x , then we let w_y take color red and w'_y take color y for every $y \neq \text{red}$. We color w with red. In the case that v_i^ℓ is colored with a color different from x , we let w_y be colored with x and w'_y be colored red for every $y \neq \text{red}$. Finally, all the neighbours of w except for v have been colored red, while v is colored with either black or white. According to the color of v , we can either color w black or white.

2. We can assume that v is red, otherwise we are done by the previous statement. For each vertex $v_i^\ell \in V_i$ and color $x \in [q] \setminus \{\mu_i(v_i^\ell)\}$, we do the following.

- If x is red, then in the construction of \widehat{C}_j we introduced a vertex w_y with list $\{y, \text{red}\}$ for every color $y \neq \text{red}$ adjacent to v_i^ℓ , and a vertex w with list $[q]$ adjacent to w_y for every $y \neq \text{red}$. Since v_i^ℓ is not colored red by μ_i , we can color w_y red for every y . Then all the neighbours of w including v have been colored with red and it is safe to color w with white.

- If x is not red, then in the construction of \widehat{C}_j we introduced two vertices w_y and w'_y for each color y except for red, and also introduced a vertex w . The vertices w_y are adjacent to v_i^ℓ and for every $y \neq \text{red}$ the vertex w'_y is adjacent to w_y . Finally, w is adjacent to all the vertices w'_y and to v . For every y , the list of w_y is $\{x, \text{red}\}$ while the list of w'_y is $\{y, \text{red}\}$. The list of w is $[q]$. Since μ_i colors v_i^ℓ with a color different from x , we let w_y be colored with x and w'_y be colored red for every $y \neq \text{red}$. Finally, all the neighbours of w including v have been colored red so it is safe to color w white.

3. Suppose for contradiction that v is red, but some vertex $v_i^\ell \in V_i$ has been colored with a color $x \neq \mu_i(v_i^\ell)$. There are two cases. If x is red, then in the construction we introduced vertices w_y adjacent to v_i^ℓ for every color $y \neq \text{red}$. Also we introduced a vertex w adjacent to v and to w_y for each $y \neq \text{red}$. The list of w_y is $\{\text{red}, y\}$ and hence w_y must have been colored y for every $y \neq \text{red}$. But then w is adjacent to v which is colored red, and to w_y which is colored y for every $y \neq \text{red}$. Thus vertex w has all colors in its neighborhood, a contradiction. In the case when x is not red, then in the construction we introduced two vertices w_y and w'_y for each $y \neq \text{red}$. Each w_y was adjacent to v_i^ℓ and had $\{x, \text{red}\}$ as its list. Since v_i^ℓ is colored x , all the w_y vertices must be colored red. For every $y \neq \text{red}$, we have that w'_y is adjacent to w_y and has $\{\text{red}, y\}$ as its list. Hence for every $y \neq \text{red}$, the vertex w'_y is colored with y . But, in the construction we also introduced a vertex w adjacent to v and to w'_y for each $y \neq \text{red}$. Thus again, vertex w has all colors in its neighbourhood, a contradiction. \square

Lemma 12. *If ϕ is satisfiable, then G has a proper list-coloring.*

Proof. Starting from a satisfying assignment of ϕ , we construct a coloring γ of G . The assignment to ϕ corresponds to a group assignment to each group F_i . Each group assignment corresponds to a coloring of V_i . For every i , we let γ color the vertices of V_i using the coloring corresponding to the group assignment of F_i .

Now we show how to complete this coloring to a proper coloring of G . Since the gadgets \widehat{C}_j are pairwise disjoint, and there are no edges going between them, it is sufficient to show that we can complete the coloring for every gadget \widehat{C}_j . Consider the clause C_j . The clause contains a literal that is set to true, and this literal belongs to a variable in some group F_i . The group assignment of F_i satisfies the clause C_j . Thus, there is a vertex v on \widehat{P}_j that corresponds to this assignment. We set $\gamma(v)$ as red (that is, γ colors v red), p_j^{start} is colored white and p_j^{end} is colored with its only admissible color, namely black if $|V(\widehat{P}_j)|$ is even and white if $|V(\widehat{P}_j)|$ is odd. The remaining vertices of \widehat{P}_j are colored alternatingly white or black. By Lemma 11(2), the coloring can be extended to every vertex of the connector between V_i and v : the coloring appearing on V_i is the coloring μ_i corresponding to the group assignment F_i . For every other vertex u on \widehat{P}_j , the color of u is black or white, thus Lemma 11(1) ensures that the coloring can be extended to any connector on u .

As this procedure can be repeated to color the gadget \widehat{C}_j for every clause C_j , we can complete γ to a proper list-coloring of G . \square

Lemma 13. *If G has a proper list-coloring γ , then ϕ is satisfiable.*

Proof. Given γ , we construct an assignment to the variables of ϕ as follows. For every group F_i of variables, if γ colors V_i with a coloring that corresponds to a group assignment of F_i , then we set this assignment for the variables in F_i . Otherwise, we set all the variables in F_i to false. We need to argue that this assignment satisfies all the clauses of ϕ .

Consider a clause C_j and the corresponding gadget \widehat{C}_j . By a simple parity argument, \widehat{P}_j cannot be colored using only the colors black and white. Thus, some vertex v on \widehat{P}_j is colored red. The vertex v corresponds to a group assignment of some group F_i that satisfies \widehat{C}_j . As

v is red, Lemma 11(3) implies that V_i is colored with the coloring μ_i that corresponds to this assignment. The construction then implies that our chosen assignment satisfies C_j . As this is true for every clause, this concludes the proof. \square

Observation 1. *The vertices $\bigcup_{i \leq t} V_i$ form a feedback vertex set of G . Furthermore, $\mathbf{pw}(G) \leq pt + 4$*

Proof. Observe that after removing $\bigcup_{i \leq t} V_i$, all that is left are the gadgets \widehat{C}_j , which do not have any edges between each other. Each such gadget is a tree and hence $\bigcup_{i \leq t} V_i$ form a feedback vertex set of G . If we place a searcher on each vertex of $\bigcup_{i \leq t} V_i$ it is easy to see that each gadget \widehat{C}_j can be searched with 4 searchers. The pathwidth bound on G follows using Proposition 1. \square

Lemma 14. *If q -LIST COLORING can be solved in $\mathcal{O}^*((q - \epsilon)^{\mathbf{fvs}(G)})$ or $\mathcal{O}^*((q - \epsilon)^{\mathbf{pw}(G)})$ time for some $\epsilon > 0$, then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Proof. Let $\mathcal{O}^*((q - \epsilon)^{\mathbf{fvs}(G)}) = \mathcal{O}^*(q^{\lambda \mathbf{fvs}(G)})$ time, where $\lambda = \log_q(q - \epsilon) < 1$. We choose a sufficiently large p such that $\delta' = \lambda \frac{p}{p-1} < 1$. Given an instance ϕ of SAT, we construct a graph G using the construction above, and run the assumed q -LIST COLORING. Correctness follows from Lemmata 12 and 13. By Observation 1, the graph G has a feedback vertex set of size $p \lceil \frac{n}{p \log q} \rceil$. The choice of p implies that

$$\lambda p \lceil \frac{n}{p \log q} \rceil \leq \lambda p \frac{n}{(p-1) \log q} + p \leq \delta' \frac{n}{\log q} + p \leq \delta'' n,$$

for some $\delta'' < 1$. Hence SAT can be solved in time $\mathcal{O}^*(2^{\delta'' n}) = \mathcal{O}^*((2 - \delta)^n)$, for some $\delta > 0$. By Observation 1, we also know that $\mathbf{pw}(G) \leq pt + 4$. Thus, the feedback vertex set size and the pathwidth of the constructed graph just differs by 4. This implies that q -LIST COLORING cannot be solved in $\mathcal{O}^*((q - \epsilon)^{\mathbf{pw}(G)})$ time. \square

Finally, observe that we can reduce q -LIST-COLORING to q -COLORING by adding a clique $Q = \{q_1, \dots, q_c\}$ on q vertices to G and making q_i adjacent to v when $i \notin L(v)$. Any coloring of Q must use q different colors, and without loss of generality q_i is colored with color i . Then one can complete the coloring if and only if one can properly color G using a color from $L(v)$ for each v . We can add the clique Q to the feedback vertex set—this increases the size of the minimum feedback vertex set by q . Since q is a constant independent of the input, this yields Theorem 4.

7 Odd Cycle Transversal

An equivalent formulation of the MAX CUT problem is to ask for a bipartite subgraph with the *maximum* number of edges, which is the same as asking for a set of edges of *minimum* size whose deletion makes the graph bipartite. We can also consider the vertex-deletion version of this problem. An *odd cycle transversal* of a graph G is a subset $S \subseteq V(G)$ such that $G \setminus S$ is bipartite. In the ODD CYCLE TRANSVERSAL problem, we are given a graph G together with an integer k and asked whether G has an odd cycle transversal of size k .

Theorem 5. *If ODD CYCLE TRANSVERSAL can be solved in $\mathcal{O}^*((3 - \epsilon)^{\mathbf{pw}(G)})$ time for $\epsilon > 0$, then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Construction. Given $\epsilon > 0$ and an instance ϕ of SAT, we construct a graph G as follows. We choose an integer p based just on ϵ . Exactly how p is chosen will be discussed at the end of this section. We start by grouping the variables of ϕ into t groups F_1, \dots, F_t of size at most $h = \lceil \log 3^p \rceil$. Thus $t = \lceil \frac{n}{\lceil \log 3^p \rceil} \rceil$. We will call an assignment of truth values to the variables in




Figure 7: Reduction to ODD CYCLE TRANSVERSAL. The arrow $A(u, v)$ from u to v with the passive odd cycle transversal shown in white (left) and the active odd cycle transversal of $A(u, v) \setminus \{u\}$ (right).

a group F_i a *group assignment*. We will say that a group assignment satisfies a clause C_j of ϕ if C_j contains at least one literal that is set to true by the group assignment. Notice that C_j can be satisfied by a group assignment of a group F_i even though C_j also contains variables that are not in F_i .

Now we describe an auxiliary gadget which will be very useful in our construction (see Figure 7). For two vertices u and v by *adding an arrow* from u to v we will mean adding a path $ua_1a_2a_3v$ on four edges starting in u and ending in v . Furthermore, we add four vertices b_1, b_2, b_3 and b_4 and edges $ub_1, b_1a_1, a_1b_2, b_2a_2, a_2b_3, b_3a_3, a_3b_4, b_4v$, and b_4v . Denote the resulting graph $A(u, v)$. None of the vertices in $A(u, v)$ except for u and v will receive any further neighbours throughout the construction of G . The graph $A(u, v)$ has the following properties, which are useful for our construction.

- The unique smallest odd cycle transversal of $A(u, v)$ is $\{a_1, a_3\}$. We call this the *passive* odd cycle transversal of the arrow.
- In $A(u, v) \setminus \{a_1, a_3\}$, u and v are in different connected components.
- The set $\{a_2, v\}$ is a smallest odd cycle transversal of $A(u, v) \setminus \{u\}$. We call this the *active* odd cycle transversal of the arrow.

The intuition behind an arrow from u to v is that if u is put into the odd cycle transversal, then v can be put into the odd cycle transversal “for free.” When the active odd cycle transversal of the arrow is picked, we say the arrow is active, otherwise we say the arrow is passive.

To construct G , we make $t \cdot p$ paths, $\{P_{i,j}\}$ for $1 \leq i \leq t, 1 \leq j \leq p$ (see Figure 8). Each path has $3m(tp+1)$ vertices, and the vertices of $P_{i,j}$ are denoted by $p_{i,j}^\ell$ for $1 \leq \ell \leq 3m(tp+1)$. For a fixed i , the paths $\{P_{i,j} : 1 \leq j \leq p\}$ correspond to the set F_i of variables. For every $1 \leq i \leq t, 1 \leq j \leq p$ and $1 \leq \ell < 3m(tp+1)$ we add three vertices $a_{i,j}^\ell, b_{i,j}^\ell$ and $q_{i,j}^\ell$ adjacent to each other. We also add the edges $a_{i,j}^\ell p_{i,j}^\ell$ and $b_{i,j}^\ell p_{i,j}^{\ell+1}$. One can think of the vertices of the paths $\{P_{i,j}\}$ layed out as rows in a matrix, where for every fixed $1 \leq \ell \leq 3m(tp+1)$ there is a column $\{p_{i,j}^\ell : 1 \leq i \leq t, 1 \leq j \leq p\}$. We group the columns three by three. In particular, For every $i \leq t$ and $0 \leq \ell < m(tp+1)$ we define the sets $P_i^\ell = \{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+2}, p_{i,j}^{3\ell+3} : 1 \leq j \leq p\}$, $A_i^\ell = \{a_{i,j}^{3\ell+1}, a_{i,j}^{3\ell+2}, a_{i,j}^{3\ell+3} : 1 \leq j \leq p\}$, $B_i^\ell = \{b_{i,j}^{3\ell+1}, b_{i,j}^{3\ell+2}, b_{i,j}^{3\ell+3} : 1 \leq j \leq p\}$ and $Q_i^\ell = \{q_{i,j}^{3\ell+1}, q_{i,j}^{3\ell+2}, q_{i,j}^{3\ell+3} : 1 \leq j \leq p\}$.

For every $i \leq t$ and $0 \leq \ell < m(tp+1)$ we make two new sets L_i^ℓ and R_i^ℓ of new vertices. Both L_i^ℓ and R_i^ℓ are independent sets of size $5p$, and we add all the edges possible between L_i^ℓ and R_i^ℓ . From L_i^ℓ we pick a special vertex λ_i^ℓ and from R_i^ℓ we pick ρ_i^ℓ . We make all the vertices in A_i^ℓ adjacent to all vertices of L_i^ℓ , and we make all vertices in B_i^ℓ adjacent to all vertices of R_i^ℓ . We make λ_i^ℓ adjacent to $\rho_i^{\ell+1}$, except for $\ell = m(tp+1) - 1$.

We will say that a subset S of P_i^ℓ which picks exactly one vertex from $P_{i,j}$ for every $1 \leq j \leq p$ is *good*. The idea is that there are $3^p \geq 2^h$ good subsets of P_i^ℓ , so we can make group assignments




Figure 8: Reduction to ODD CYCLE TRANSVERSAL. The path $P_{i,j}$ with three different ways of removing a set Z and partitioning the remaining bipartite graph into classes L and R .

of F_i correspond to good subsets of P_i^ℓ . For every good subset S of P_i^ℓ we add a cycle $X_{i,S}^\ell$. The cycle $X_{i,S}^\ell$ has length $2p + 1$. We select a vertex on $X_{i,S}^\ell$ and call it $x_{i,S}^\ell$. For every vertex $u \in P_i^\ell \setminus S$ we add an arrow from u to a vertex of $X_{i,S}^\ell$. We add arrows in such a way that every vertex of $X_{i,S}^\ell \setminus \{x_{i,S}^\ell\}$ is the endpoint of exactly one arrow.

For every $i \leq t$ and $0 \leq \ell < m(tp+1)$, we make a cycle Y_i^ℓ of length 3^p . Notice that the length of the cycle is odd. Every vertex of Y_i^ℓ corresponds to a good subset S of P_i^ℓ . For each good subset S of P_i^ℓ we add an arrow from $x_{i,S}^\ell$ of the cycle $X_{i,S}^\ell$ to the vertex in Y_i^ℓ that corresponds to S .

We say that a good subset of P_i^ℓ is *equal* with a good subset S' of $P_i^{\ell'}$ if for every $1 \leq j \leq t$, the distance along $P_{i,j}$ between the vertex of S on $P_{i,j}$ and the vertex of S' on $P_{i,j}$ is divisible by 3. Informally, S and S' are equal if they look identical when we superimpose P_i^ℓ onto $P_i^{\ell'}$. To every group assignment of variables F_i , we designate a good subset of P_i^ℓ for every ℓ . We designate good subsets in such a way that good subsets corresponding to the same group assignment are equal.

Finally, for every clause C_j , $1 \leq j \leq m$, we will introduce $tp + 1$ cycles. That is, for every $0 \leq r \leq tp$, we introduce a cycle \widehat{C}_j^r . The cycle contains one vertex for every $i \leq t$ and group assignment to F_i , and potentially one dummy vertex to make it have odd length. Going around the cycle counterclockwise we first encounter all the vertices corresponding to group assignments of F_1 , then all the vertices corresponding to group assignments of F_2 , and so on. For $i \leq t$ and every good subset S of P_i^{rm+j} that corresponds to a group assignment of F_i that satisfies C_j we add an arrow from $x_{i,S}^{rm+j}$ to the vertex on \widehat{C}_j^r that corresponds to the same group assignment of F_i as S does. This concludes the construction of G .

The intention behind the construction is that if ϕ is satisfiable, then a minimum odd cycle transversal of G can pick:

- One vertex from each triangle $\{a_{i,j}^\ell, b_{i,j}^\ell, q_{i,j}^\ell\}$ for each $1 \leq i \leq t$, $1 \leq j \leq p$, $1 \leq \ell < 3m(tp + 1)$. There are $tp(3m(tp + 1) - 1)$ such triangles in total.
- One vertex from $\{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+2}, p_{i,j}^{3\ell+3}\}$ for each $1 \leq i \leq t$, $1 \leq j \leq p$, $0 \leq \ell < m(tp + 1)$. There are $tpm(tp + 1)$ such triples.
- Two vertices from every arrow added, *without* counting the starting point of the arrow. For each $i \leq t$ and $0 \leq \ell < m(tp + 1)$, there are $2p3^p$ arrows ending in some cycle $X_{i,S}^\ell$. Hence there are $2p3^p tm(tp + 1)$ such arrows. For every $i \leq t$ and $0 \leq \ell < m(tp + 1)$ there are 3^p arrows ending in the cycle Y_i^ℓ . Hence there are $3^p tm(tp + 1)$ such arrows. For every clause C_j , there are $tp + 1$ arrows added for every group assignment that satisfies that clause. Let μ be the sum over all clauses of the number of group assignments that satisfy that clause. The total number of arrows added is then $\mu(tp + 1) + (2p + 1)3^p tm(tp + 1)$. Thus the odd cycle transversal can pick $2\mu(tp + 1) + 2(2p + 1)3^p tm(tp + 1)$ vertices from arrows.
- One vertex $x_{i,S}^\ell$ for every $i \leq t$ and $0 \leq \ell < m(tp + 1)$. There are $tm(tp + 1)$ choices for i and ℓ .

We let the α be the value of the total budget, that is the sum of the items above.

Lemma 15. *If ϕ is satisfiable, then G has an odd cycle transversal of size α .*

Proof. Given a satisfying assignment γ to ϕ , we construct an odd cycle transversal Z of G of size α together with a partition of $V(G) \setminus Z$ into L and R such that every edge of $G \setminus Z$ goes between a vertex in L and a vertex in R . The assignment to ϕ corresponds to a group assignment of each F_i for $1 \leq i \leq t$. For every $1 \leq i \leq t$ and $0 \leq \ell < m(tp + 1)$, we add to Z the good subset S of P_i^ℓ that corresponds to the group assignment of F_i . Notice that for each fixed i , the sets

picked from P_i^ℓ and $P_i^{\ell'}$ are equal for any ℓ, ℓ' . At this point we have picked one vertex from $\{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+2}, p_{i,j}^{3\ell+3}\}$ for each $1 \leq i \leq t, 1 \leq j \leq p, 0 \leq \ell < m(tp+1)$.

For every fixed $1 \leq i \leq t, 1 \leq j \leq p$, there are three cases. If $p_{i,j}^1 \in Z$, we put $p_{i,j}^2$ into L and $p_{i,j}^3$ into R . If $p_{i,j}^2 \in Z$ we put $p_{i,j}^1$ into R and $p_{i,j}^3$ into L . If $p_{i,j}^3 \in Z$ we put $p_{i,j}^1$ into L and $p_{i,j}^2$ into R . Now, for every $4 \leq \ell \leq 3m(tp+1)$ such that $p_{i,j}^\ell \notin Z$ we put $p_{i,j}^\ell$ into the same set out of $\{L, R\}$ as $p_{i,j}^{\ell'}$ where $1 \leq \ell' \leq 3$ and $\ell \equiv \ell' \pmod{3}$.

For every $1 \leq i \leq t, 0 \leq \ell \leq m(tp+1)$, we put L_i^ℓ into L and R_i^ℓ into R . For every triple of a, b, q of pairwise adjacent vertices such that $a \in A_i^\ell, b \in B_i^\ell$, and $q \in Q_i^\ell$, we proceed as follows. The vertex a has a neighbour a' in P_i^ℓ and b has a neighbour b' in P_i^ℓ . There is a j such that b' is the successor of a' on $P_{i,j}$. Thus, there are three cases;

- $a' \in Z$ and $b' \in L$, we put a in R , q in L and b in Z .
- $a' \in R$ and $b' \in Z$, we put a in Z , q in R and b in L .
- $a' \in L$ and $b' \in R$, we put a in R , q in Z and b in L .

For every $1 \leq i \leq t, 0 \leq \ell \leq m(tp+1)$, there are many arrows from vertices in P_i^ℓ to vertices on cycles $X_{i,S}^\ell$ for good subsets S of P_i^ℓ . For each arrow, if its endpoint in P_i^ℓ is in Z we add the active odd cycle transversal of the arrow to Z , otherwise we add the passive odd cycle transversal of the arrow to Z . In either case, the remaining vertices on the arrow form a forest, and therefore we can insert the remaining vertices of the arrow into L and R according to which sets out of $\{L, R, Z\}$ u and v are in.

For every $1 \leq i \leq t, 0 \leq \ell \leq m(tp+1)$, there is exactly one set S such that the cycle $X_{i,S}^\ell$ only has passive arrows pointing into it. This is exactly the set S which corresponds to the restriction of γ to F_i . Each cycle $X_{i,S'}^\ell$ that has at least one arrow pointing into them already contains at least one vertex in Z —the endpoint of the active arrow pointing into the cycle. Thus we can partition the remaining vertices of $X_{i,S'}^\ell$ into L and R such that no edge has both endpoints in L or both endpoints in R . For the cycle $X_{i,S}^\ell$, we put $x_{i,S}^\ell$ into Z and partition the remaining vertices of $X_{i,S}^\ell$ into L and R such that no edge has both endpoints in L or both endpoints in R . We add the active odd cycle transversal in the arrow from $x_{i,S}^\ell$ to the cycle Y_i^ℓ into Z . For all other good subsets S' , we add the passive odd cycle transversal in the arrow from $x_{i,S}^\ell$ to the cycle Y_i^ℓ into Z . Thus each cycle Y_i^ℓ contains one vertex in Z and the remaining vertices of Y_i^ℓ can be distributed into L and R .

For every arrow that goes from a vertex $x_{i,S}^\ell$ into a cycle \widehat{C}_h^r , we add the active odd cycle transversal of the arrow to Z if $x_{i,S}^\ell \in Z$ and add the passive odd cycle transversal to Z otherwise. Again the remaining vertices on each arrow can easily be partitioned into L and R such that no edge has both endpoints in L or both endpoints in R . This concludes the construction of Z . Since we have put the vertices into Z in accordance to the budget described in the construction it follows that $|Z| \leq \alpha$. All that remains to show, is that for each $1 \leq h \leq m$ and $0 \leq r \leq tp$, the cycle \widehat{C}_h^r has at least one active arrow pointing into it.

The cycle \widehat{C}_h^r corresponds to the clause C_h . The clause C_h is satisfied by γ and hence it is satisfied by the restriction of γ to some group F_i . This restriction is a group assignment of F_i and hence it corresponds to a good subset S of P_i^{rm+h} , which happens to be exactly $Z \cap P_i^{rm+h}$. Thus $x_{i,S}^{rm+h} \in Z$ and since the restriction of γ to F_i satisfies C_h , there is an arrow pointing from $x_{i,S}^{rm+h}$ and into \widehat{C}_h^r . Since this arrow is active, this concludes the proof. \square

Lemma 16. *If G has an odd cycle transversal of size α , then ϕ is satisfiable.*

Proof. Let Z be an odd cycle transversal of G of size α . Since $G \setminus Z$ is bipartite, the vertices of $G \setminus Z$ can be partitioned into L and R such that every edge of $G \setminus Z$ has one endpoint in L and the other in R . Given Z, L and R , we construct a satisfying assignment to ϕ . Every arrow

in G must contain at least two vertices in Z , not counting the startpoint of the arrow. Let \vec{Z} be a subset of Z containing two vertices from each arrow, but no arrow start point. Observe that no two arrows have the same endpoint, and therefore $|\vec{Z}|$ is exactly two times the number of arrows in G . Let $Z' = Z \setminus \vec{Z}$.

We argue that for any $1 \leq i \leq t$ and $0 \leq \ell < m(tp + 1)$ we have $|Z' \cap (L_i^\ell \cup R_i^\ell \cup A_i^\ell \cup B_i^\ell \cup Q_i^\ell \cup P_i^\ell)| \geq 4p$. Observe that no vertices in $L_i^\ell, R_i^\ell, A_i^\ell, B_i^\ell, Q_i^\ell$ or P_i^ℓ are endpoints of arrows, and hence they do not contain any vertices of \vec{Z} . Suppose for contradiction that $|Z' \cap (L_i^\ell \cup R_i^\ell \cup A_i^\ell \cup B_i^\ell \cup Q_i^\ell \cup P_i^\ell)| < 4p$. Then there is a vertex $\lambda \in L_i^\ell \setminus Z'$, and a vertex $\rho \in R_i^\ell \setminus Z'$. Without loss of generality, $\lambda \in L$ and $\rho \in R$. Furthermore, there is a $1 \leq j \leq p$ such that

$$|Z' \cap \{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+2}, p_{i,j}^{3\ell+3}, a_{i,j}^{3\ell+1}, a_{i,j}^{3\ell+2}, a_{i,j}^{3\ell+3}, b_{i,j}^{3\ell+1}, b_{i,j}^{3\ell+2}, b_{i,j}^{3\ell+3}, q_{i,j}^{3\ell+1}, q_{i,j}^{3\ell+2}, q_{i,j}^{3\ell+3}\}| < 4.$$

Since $\{a_{i,j}^{3\ell+1}, b_{i,j}^{3\ell+1}, q_{i,j}^{3\ell+1}\}$, $\{a_{i,j}^{3\ell+2}, b_{i,j}^{3\ell+2}, q_{i,j}^{3\ell+2}\}$ and $\{a_{i,j}^{3\ell+3}, b_{i,j}^{3\ell+3}, q_{i,j}^{3\ell+3}\}$ form triangles and must contain a vertex from Z' each, it follows that each of these triangles contain exactly one vertex from Z' , and that $Z' \cap \{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+2}, p_{i,j}^{3\ell+3}\} = \emptyset$. Since $\lambda \in L$ and $\rho \in R$, λ is adjacent to all vertices of $A_{i,j}^\ell$ and ρ is adjacent to all vertices of $B_{i,j}^\ell$, it follows that $A_{i,j}^\ell \setminus Z' \subseteq R$ and $B_{i,j}^\ell \setminus Z' \subseteq L$.

Hence, there are two cases to consider either (1) $\{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+3}\} \subseteq L$ and $p_{i,j}^{3\ell+2} \in R$ or (2) $\{p_{i,j}^{3\ell+1}, p_{i,j}^{3\ell+3}\} \subseteq R$ and $p_{i,j}^{3\ell+2} \in L$. In the first case, observe that either $a_{i,j}^{3\ell+2} \in R$ or $b_{i,j}^{3\ell+2} \in L$ and hence either $a_{i,j}^{3\ell+2} p_{i,j}^{3\ell+2}$ or $b_{i,j}^{3\ell+2} p_{i,j}^{3\ell+3}$ have both endpoints in the same set out of $\{L, R\}$, a contradiction. The second case is similar, either $a_{i,j}^{3\ell+1} \in R$ or $b_{i,j}^{3\ell+1} \in L$ and hence either $a_{i,j}^{3\ell+1} p_{i,j}^{3\ell+1}$ or $b_{i,j}^{3\ell+1} p_{i,j}^{3\ell+2}$ have both endpoints in the same set out of $\{L, R\}$, a contradiction. We conclude that $|Z' \cap (L_i^\ell \cup R_i^\ell \cup A_i^\ell \cup B_i^\ell \cup Q_i^\ell \cup P_i^\ell)| \geq 4p$.

For any $1 \leq i \leq t$ and $0 \leq \ell < m(tp + 1)$, Y_i^ℓ is an odd cycle so Y_i^ℓ contains a vertex in Z . If Y_i^ℓ contains no vertices of Z' , then it contains a vertex from \vec{Z} and there is an active arrow pointing into Y_i^ℓ . The starting point of this arrow is a vertex $x_{i,S}^\ell$ for some good subset S of P_i^ℓ . Since the arrow is active and $x_{i,S}^\ell$ is not the endpoint of any arrow, we know that $x_{i,S}^\ell \in Z'$. Hence for any $1 \leq i \leq t$ and $0 \leq \ell < m(tp + 1)$, we have that either there is a good subset S of P_i^ℓ such that $x_{i,S}^\ell \in Z'$ or at least one vertex of Y_i^ℓ is in Z' .

The above arguments, together with the budget constraints, imply that for every $1 \leq i \leq t$ and $0 \leq \ell < m(tp + 1)$, we have $|Z' \cap (L_i^\ell \cup R_i^\ell \cup A_i^\ell \cup B_i^\ell \cup Q_i^\ell \cup P_i^\ell)| = 4p$ and that $|Z' \cap \bigcup \{x_{i,S}^\ell\} \cup V(Y_i^\ell)| = 1$, where the union is taken over all good subsets S of P_i^ℓ . It follows $Z' \cap P_i^\ell$ is a good subset of P_i^ℓ . Let $S = Z' \cap P_i^\ell$. The cycle $X_{i,S}^\ell$ has odd length, and hence it must contain some vertex from Z . On the other hand, all the arrows pointing into $X_{i,S}^\ell$ are passive, so $X_{i,S}^\ell$ cannot contain any vertices from \vec{Z} . Thus $X_{i,S}^\ell$ contains a vertex from Z' , and by the budget constraints this must be $x_{i,S}^\ell$.

Now, consider three consecutive vertices $p_{i,j}^\ell, p_{i,j}^{\ell+1}, p_{i,j}^{\ell+2}$ for some $1 \leq i \leq t, 1 \leq j \leq p, 1 \leq \ell \leq 3m(tp + 1) - 2$. We prove that at least one of them has to be in Z . Suppose not. We know that neither $\lambda_i^{[\ell/3]}, \rho_i^{[\ell/3]}, \lambda_i^{[\ell/3]+1}$ nor $\rho_i^{[\ell/3]+1}$ are in Z . Thus, without loss of generality $\{\lambda_i^{[\ell/3]}, \lambda_i^{[\ell/3]+1}\} \subseteq L$ and $\{\rho_i^{[\ell/3]}, \rho_i^{[\ell/3]+1}\} \subseteq R$. There are two cases. Either $p_{i,j}^\ell \in R$ and $p_{i,j}^{\ell+1} \in L$ or $p_{i,j}^{\ell+1} \in L$ and $p_{i,j}^{\ell+2} \in R$. In the first case, we obtain a contradiction since either $a_{i,j}^\ell \in R$ or $b_{i,j}^\ell \in L$. In the second case, we get a contradiction since either $a_{i,j}^{\ell+1} \in R$ or $b_{i,j}^{\ell+1} \in L$. Hence for any three consecutive vertices on $P_{i,j}$, at least one of them is in Z . Since the budget constraints ensure that there are at most $|V(P_{i,j})|/3$ vertices in $P_{i,j} \cap Z$, it follows from the pigeon hole principle that there is an $0 \leq r \leq tp$ such that for any $1 \leq i \leq t$ and $1 \leq h \leq m$ and

$1 \leq h' \leq m$, the set $P_i^{rm+h} \cap Z$ equals $P_i^{rm+h'} \cap Z$. Here equality is in the sense of equality of good subsets of P_i^ℓ .

For every $1 \leq i \leq t$, $P_i^{rm+1} \cap Z$ is a good subset of P_i^{rm+1} . If $P_i^{rm+1} \cap Z$ corresponds to a group assignment of F_i , then we set the variables in F_i to this assignment. Otherwise we set all the variables in F_i to false. We need to argue that every clause C_h is satisfied by this assignment. Consider the cycle \widehat{C}_h^r . Since it is an odd cycle, it must contain a vertex from Z , the budget constraints and the discussion above implies that this vertex is from \vec{Z} . Hence there must be an active arrow pointing into \widehat{C}_h^r . The starting point of this active arrow is a vertex $x_{i,S}^{mr+h}$ for some i and good subset S of P_i^{mr+h} . The set S corresponds to a group assignment of F_i that satisfies C_h . Since the arrow is active $x_{i,S}^{mr+h} \in Z'$, and by the discussion above we have that $P_i^{mr+h} \cap Z' = S$. Now, $S = P_i^{mr+h} \cap Z'$ and S is equal to $P_i^{mr+1} \cap Z'$ and hence the assignment to the variables of F_i satisfies C_h . Since this holds for all clauses, this concludes the proof. \square

Lemma 17. $\text{pw}(G) \leq t(p+1) + 10p3^p$.

Proof. We show how to search the graph using at most $t(p+1) + 10p3^p$ searchers. The strategy consists of $m(tp+1)$ rounds numbered from round 0 to round $m(tp+1) - 1$. Each round has t stages, numbered from 1 to t . In the beginning of round k there is a searcher on $p_{i,j}^{3k+1}$ and ρ_i^k for every $1 \leq i \leq t$, $1 \leq j \leq p$. Let r and $1 \leq h \leq m$ be integers such that $k+1 = rm+h$. Recall, that as we go around \widehat{C}_h^r counterclockwise we first encounter vertices corresponding to group assignments of F_1 , then to assignments of F_2 and so on. In the beginning of round k we place a searcher on the first vertex on \widehat{C}_h^r that corresponds to an assignment of F_1 . If \widehat{C}_h^r contains a dummy vertex, we place a searcher on this vertex as well. These two searchers will remain on their respective vertices throughout the round. In the beginning of stage s of round k we will assume that the vertices on the cycle \widehat{C}_h^r corresponding to group assignments of $F_{s'}$, $s' < s$ have already been cleaned, and in the beginning of every stage $s > 1$, there is a searcher standing on the first vertex corresponding to a group assignment of F_s .

In stage s of round k , we place searchers on all vertices of $P_s^k, A_s^k, B_s^k, Q_s^k, L_s^k, R_s^k, Y_s^k$ and all vertices of cycles $X_{s,S}^k$ for every good subset S of P_s^k , on all vertices of arrows starting or ending in such cycles, and on all vertices of \widehat{C}_h^r corresponding to group assignments of F_s . In total this amounts to less than $10p3^p$ vertices.

In the last part of stage s of round k , we place searchers on $p_{s,j}^{3(k+1)+1}$ for every $1 \leq j \leq p$ and on ρ_s^{k+1} . Then we remove all the searchers that were placed out in the first part of phase s except for the searcher on the last vertex on \widehat{C}_h^r corresponding to a group assignment of F_s . Unless $s = 1$ there is also a searcher on the last vertex on \widehat{C}_h^r corresponding to a group assignment of F_{s-1} . We remove this searcher, and the next stage can commence. In the end of the last stage of round k we remove all the searchers from \widehat{C}_h^r . Then the last stage can commence. At any point in time, at most $t(p+1) + 10p3^p$ searchers are placed on G . \square

Proof (of Theorem 5). Suppose ODD CYCLE TRANSVERSAL can be solved in time $\mathcal{O}^*((3-\epsilon)^{\text{pw}(G)})$ for some $\epsilon > 0$. Then there is an $\epsilon' < 1$ such that $\mathcal{O}^*((3-\epsilon)^{\text{pw}(G)}) \leq \mathcal{O}^*(3^{\epsilon' \text{pw}(G)})$. We choose p large enough such that $\epsilon' \cdot \frac{p+1}{p-1} = \delta' < 1$. Given an instance of SAT we construct an instance of ODD CYCLE TRANSVERSAL using the above construction and the chosen value of p . Then we solve the ODD CYCLE TRANSVERSAL instance using the $\mathcal{O}^*((3-\epsilon)^{\text{pw}(G)})$ time algorithm. Correctness is ensured by Lemmata 15 and 16. Lemma 17 yields that the total time taken is upper bounded by $\mathcal{O}^*((3-\epsilon)^{\text{pw}(G)}) \leq \mathcal{O}^*(3^{\epsilon' \text{pw}(G)}) \leq \mathcal{O}^*(3^{\epsilon'(t(p+1)+f(\epsilon'))}) \leq \mathcal{O}^*(3^{\epsilon' \lceil \frac{n}{\lfloor p \log 3 \rfloor} \rceil (p+1)}) \leq \mathcal{O}^*(3^{\epsilon' \frac{n(p+1)}{\lfloor p \log 3 \rfloor}}) \leq \mathcal{O}^*(3^{\epsilon' \frac{n(p+1)}{(p-1) \log 3}}) \leq \mathcal{O}^*(3^{\delta' \frac{n}{\log 3}}) \leq \mathcal{O}^*(2^{\delta' n}) = \mathcal{O}^*((2-\delta)^n)$ for $\delta < 1$. \square




Figure 9: Reduction to TRIANGLE PACKING, showing how the vertices \widehat{c}_j^r and \widehat{d}_j^r representing clause $C_j = (x_1 \vee \bar{x}_2 \vee x_4)$ are connected to the paths P_1, \dots, P_n .

8 Partition Into Triangles

A *triangle packing* in a graph G is a collection of pairwise disjoint vertex sets S_1, S_2, \dots, S_t in G such that S_i induces a triangle in G for every i . The size of the packing is t . If $V(G) = \bigcup_{i \leq t} S_i$, then the collection $S_1 \dots S_t$ is a *partition of G into triangles*. In the TRIANGLE PACKING problem, we are given a graph G and an integer t and asked whether there is a triangle packing in G of size at least t . In the PARTITION INTO TRIANGLES problem, we are given a graph G and asked whether G can be partitioned into triangles. Notice that since PARTITION INTO TRIANGLES is the special case of TRIANGLE PACKING when the number of triangles is the number of vertices divided by 3, the bound of Theorem 6 holds for TRIANGLE PACKING as well.

Theorem 6. *If PARTITION INTO TRIANGLES can be solved in time $\mathcal{O}^*((2 - \epsilon)^{\text{pw}(G)})$ for $\epsilon > 0$, then SAT can be solved in $\mathcal{O}^*((2 - \delta)^n)$ time for some $\delta > 0$.*

Construction. first show the lower bound for TRIANGLE PACKING and then modify our construction to also work for the more restricted PARTITION INTO TRIANGLES problem. Given an instance ϕ of SAT we construct a graph G as follows (see Figure 9). For every variable v_i , we make a path P_i on $2m(n + 1) + 1$ vertices. We denote the ℓ -th vertex of P_i by p_i^ℓ . For every i , we add a set T_i of $2m(n + 1)$ vertices, and let the ℓ -th vertex of T_i be denoted t_i^ℓ . For every $1 \leq \ell \leq 2m(n + 1)$ we add the edges $t_i^\ell p_i^\ell$ and $t_i^\ell p_i^{\ell+1}$.

For every clause C_j , we add $n + 1$ gadgets corresponding to the clause. In particular, for every $0 \leq r \leq n$ we do the following. First we add the vertices \widehat{c}_j^r and \widehat{d}_j^r and the edge $\widehat{c}_j^r \widehat{d}_j^r$. For every variable v_i that occurs in C_j positively, we add the edges $\widehat{c}_j^r t_i^{2(mr+j)}$ and $\widehat{d}_j^r t_i^{2(mr+j)}$. For every variable v_i that occurs in C_j negated, we add the edges $\widehat{c}_j^r t_i^{2(mr+j)-1}$ and $\widehat{d}_j^r t_i^{2(mr+j)-1}$. Doing this for every r and every clause C_j concludes the construction of G .

Lemma 18. *If ϕ satisfiable, then G has a triangle packing of size $mn(n + 1) + m(n + 1)$.*

Proof. Consider a satisfying assignment to ϕ . For every variable v_i that is set to true and integer $1 \leq \ell \leq m(n+1)$, we add $\{t_i^{2\ell-1}, p_i^{2\ell-1}, p_i^{2\ell}\}$ to the triangle packing. For every variable v_i that is set to false and integer $1 \leq \ell \leq m(n+1)$, we add $\{t_i^{2\ell}, p_i^{2\ell}, p_i^{2\ell+1}\}$ to the triangle packing. For every clause C_j , there is a literal set to true. Suppose this literal corresponds to the variable v_i . Notice that if v_i occurs positively in C_j , then v_i is set to true, and if it occurs negatively it is set to false. For each $0 \leq r \leq n$, if v_i occurs positively in C_j , then $t_i^{2(mr+j)}$ has not yet been used in any triangle, so we can add $\{\widehat{c}_j^r, \widehat{d}_j^r, t_i^{2(mr+j)}\}$ to the triangle packing. On the other hand, if v_i occurs negated in C_j , then $t_i^{2(mr+j)-1}$ has not yet been used in any triangle, so we can add $\{\widehat{c}_j^r, \widehat{d}_j^r, t_i^{2(mr+j)-1}\}$ to the triangle packing. In total $mn(n+1) + m(n+1)$ triangles are packed. \square

Lemma 19. *If G has a triangle packing of size $mn(n+1) + m(n+1)$, then ϕ is satisfiable.*

Proof. Observe that for any j and r , every triangle that contains \widehat{c}_j^r also contains \widehat{d}_j^r and vice versa. Furthermore, if we remove all the vertices \widehat{c}_j^r and \widehat{d}_j^r for every j and r from G we obtain a disconnected graph with n connected components, $G[T_i \cup V(P_i)]$ for every i . Thus, the only way to pack $mn(n+1) + m(n+1)$ triangles in G is to pack $mn(n+1)$ triangles in each component $G[T_i \cup V(P_i)]$ and in addition make sure that every pair $(\widehat{c}_j^r, \widehat{d}_j^r)$ is used in some triangle in the packing.

The only way to pack $mn(n+1)$ triangles in a component $G[T_i \cup V(P_i)]$ is to use every second triangle of the form $\{t_i^\ell, p_i^\ell, p_i^{\ell+1}\}$, except possibly at one point where two triangles on this form are skipped. By the pigeon hole principle there is an $0 \leq r \leq n$ such that for every i , every second triangle of the form $\{t_i^{2mr+\ell}, p_i^{2mr+\ell}, p_i^{2mr+\ell+1}\}$ for $1 \leq \ell \leq 2m$ is used. We make an assignment to the variables of ϕ as follows. For every i such that $\{t_i^{2mr+1}, p_i^{2mr+1}, p_i^{2mr+2}\}$ is used, v_i is set to true, and otherwise $\{t_i^{2mr+2}, p_i^{2mr+2}, p_i^{2mr+3}\}$ is used in the packing and v_i is set to false. We prove that this assignment satisfies ϕ .

For every j , the pair $(\widehat{c}_j^r, \widehat{d}_j^r)$ is used in some triangle in the packing. This triangle either contains $t_i^{2(mr+j)}$ or $t_i^{2(mr+j)-1}$ for some i . If it contains $t_i^{2(mr+j)}$, then v_i occurs positively in C_j . Furthermore, since the triangle packing contains every second triangle of the form $\{t_i^{2mr+\ell}, p_i^{2mr+\ell}, p_i^{2mr+\ell+1}\}$ for $1 \leq \ell \leq 2m$, it follows that the triangle packing contains $\{t_i^{2mr+1}, p_i^{2mr+1}, p_i^{2mr+2}\}$ and hence v_i is set to true. By an identical argument, if the triangle containing the pair $(\widehat{c}_j^r, \widehat{d}_j^r)$ contains $t_i^{2(mr+j)-1}$, then v_i occurs negated in C_j and v_i is set to false. This concludes the proof. \square

We now modify the construction to work for PARTITION INTO TRIANGLES instead of TRIANGLE PACKING. Given the graph G as constructed from ϕ , we construct a graph G' as follows. For every $1 \leq i \leq n$ and $1 \leq l \leq m(n+1)$, we make a clique Q_i^ℓ on four vertices. The vertices of Q_i^ℓ are all adjacent to t_i^{2l} and to t_i^{2l-1} . For every $i < n$ and $1 \leq l \leq m(n+1)$ we make all vertices of Q_i^ℓ adjacent to all vertices of Q_{i+1}^ℓ . Suppose that $2n+2$ is p modulo 3 for some $p \in \{0, 1, 2\}$. We remove p vertices from Q_n^ℓ for every $l \leq m(n+1)$.

Lemma 20. *G has a triangle packing of size α if and only if G' can be partitioned into triangles. Here, α is a non-negative integer.*

Proof. In the forward direction, consider a triangle packing of size α in G as constructed in Lemma 18. We can assume that the triangle packing has this form, because by Lemma 19 we have that ϕ is satisfiable.

For every fixed $1 \leq l \leq m(n+1)$, we proceed as follows. We know that there exists an i such that both t_i^{2l} and t_i^{2l-1} are used in the packing. For every $i' \neq i$, exactly one out of $t_{i'}^{2l}$ and $t_{i'}^{2l-1}$ is used in the packing. For each such i' , we make a triangle containing the unused vertex out of $t_{i'}^{2l}$ and $t_{i'}^{2l-1}$ and two vertices of $Q_{i'}^\ell$. Then we “clean up” $Q_1^\ell, \dots, Q_n^\ell$ as follows.

In particular, we start with the yet unused vertices of Q_1^ℓ . There are two of them. Make a triangle containing these two vertices and one vertex of Q_2^ℓ . Now Q_2^ℓ has one unused vertex left. Make a triangle containing this vertex and the two unused vertices of Q_3^ℓ . Continue in this fashion until arrive at $Q_{i'}^\ell$. At this point we have used 0, 1 or 2 vertices of $Q_{i'}^\ell$, a triangle containing some vertices in $Q_{i'-1}^\ell$. The case when we have used 0 vertices of $Q_{i'}^\ell$ also covers the case that $i' = 1$. If we only used 0 or 1 vertices of $Q_{i'}^\ell$, then we add a triangle that contains 3 vertices of $Q_{i'}^\ell$. If there are still unused vertices in $Q_{i'}^\ell$, then their number is either 1 or 2. We make a triangle containing these vertices and 1 or 2 of the unused vertices of $Q_{i'+1}^\ell$. Now we proceed to $Q_{i'+1}^\ell$ and continue in this manner until we reach Q_n^ℓ . Since the total number of vertices in $\bigcup_{j \leq n} Q_j^\ell$ is $4n - p$, we know that $2n - 2$ of these vertices are used for triangles with vertices of G , and $2n + 2 - p$ is divisible by 3 the process described above will partition all the unused vertices of $\bigcup_{j \leq n} Q_j^\ell$ into triangles.

In the reverse direction, we argue that in any partitioning of G' into triangles, exactly α triangles must lie entirely within G . In fact, we argue that for any $l \leq m(n + 1)$ exactly $n - 1$ vertices out of $\bigcup_{i \leq n} \{t_i^{2l}, t_i^{2l-1}\}$ are used in triangles containing vertices from $\bigcup_{i \leq n} Q_i^\ell$.

Pick $1 \leq j \leq m$ and r such that $l = mr + j$. Exactly one out of $\bigcup_{i \leq n} \{t_i^{2l}, t_i^{2l-1}\}$ is in a triangle with \widehat{c}_j^r and \widehat{d}_j^r . Furthermore, for each $i \leq n$ the vertex p_i^{2l} must be in a triangle either containing t_i^{2l} or t_i^{2l-1} . Hence, at most $n - 1$ vertices out of $\bigcup_{i \leq n} \{t_i^{2l}, t_i^{2l-1}\}$ are used in triangles containing vertices from $\bigcup_{i \leq n} Q_i^\ell$. Furthermore, any triangle containing t_i^{2l} or t_i^{2l-1} must either contain p_i^{2l} , \widehat{c}_j^r or some vertex in $\bigcup_{i \leq n} Q_i^\ell$. Hence exactly $n - 1$ vertices out of $\bigcup_{i \leq n} \{t_i^{2l}, t_i^{2l-1}\}$ are used in triangles containing vertices from $\bigcup_{i \leq n} Q_i^\ell$. Thus in the packing, exactly 3α vertices in G' are contained in triangles completely inside G , and hence G has a triangle packing of size α . \square

To complete the proof for PARTITION INTO TRIANGLES we need to bound the pathwidth of G' .

Lemma 21. $\text{pw}(G') \leq n + 10$.

Proof. We give a search strategy for G' that uses $n + 10$ searchers. The strategy consists of $m(n + 1)$ rounds and each round has n stages. In the beginning of round l , $1 \leq l \leq m(n + 1)$, there are n searchers placed, one on each vertex p_i^{2l-1} for every i . Let r and $1 \leq j \leq m$ be integers such that $l = mr + j$. We place one searcher on \widehat{c}_j^r and one on \widehat{d}_j^r . These two searchers will stay put throughout the duration of this round. In stage i of round l we place searchers on all vertices of Q_i^ℓ and Q_{i+1}^ℓ . Then we place searchers on t_i^{2l-1} , t_i^{2l} , p_i^{2l} and p_i^{2l+1} . At the end of stage i we remove the searchers from Q_i^ℓ , t_i^{2l-1} , t_i^{2l} and p_i^{2l} . We then proceed to the next stage. At the end of the round we remove the searchers from \widehat{c}_j^r and \widehat{d}_j^r . Notice that now, there are searchers on p_i^{2l+1} for every i , and the next round can commence. \square

Lemmata 18,19,20 and 21 prove Theorem 6.

9 Conclusion

We have showed that for a number of basic graph problems, the best known algorithms parameterized by treewidth are optimal in the sense that base of the exponential dependence on treewidth is best possible. Recall that for DOMINATING SET and PARTITION INTO TRIANGLES, this running time was obtained using the technique of fast subset sum convolutions [50]. Thus it could have been a real possibility that the running time is improved for some other problems as well.

The results are proved under the Strong Exponential Time Hypothesis (SETH). While this hypothesis is extremely strong and might not be accepted by everyone, our results at least make a connection between rather specific graph problems and the very basic issue of better SAT algorithms. Our results suggest that one should not try to find better algorithms on bounded treewidth graphs for the problems considered in the paper: as this would disprove SETH, such an effort is better spent on trying to disprove SETH directly in the domain of satisfiability. Finally, we suggest the following open questions for future work:

- Can we prove similar tight lower bounds under the restriction that the graph is planar? Or is it possible to find improved algorithms on bounded treewidth planar graphs?
- For the q -COLORING problem, we were able to prove lower bounds parameterized by the feedback vertex set number. Can we prove such bounds for the other problems as well? Recently, Jaffke and Jansen [33] strengthened our lower bounds for q -COLORING. In particular, they showed that q -COLORING parameterized by the modulator to linear forests (a forest where every connected component is a path), say $\text{lfvs}(G)$, can not be solved in time $(q - \epsilon)^{\text{lfvs}(G)} |V(G)|^{\mathcal{O}(1)}$.

Acknowledgements. We sincerely thank all the reviewers for their insightful comments and suggestions. Daniel Lokshtanov was supported by the Bergen Research Foundation and the University of Bergen through project “BeHard”. Furthermore, he was supported by ERC Starting Grant PaPaAlg (No. 715744). Saket Saurabh was supported by the ERC Starting Grant PARAPPROX (No. 306992). Dániel Marx was supported by ERC Starting Grant PARAMTIGHT (No. 280152) and Consolidator Grant SYSTEMATICGRAPH (No. 755978).

References

- [1] Amir Abboud, Arturs Backurs, and Virginia Vassilevska Williams. Tight hardness results for LCS and other sequence similarity measures. In *Proceedings of the 56th Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pages 59–78, 2015.
- [2] Amir Abboud and Virginia Vassilevska Williams. Popular conjectures imply strong lower bounds for dynamic problems. In *Proceedings of the 55th Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pages 434–443, 2014.
- [3] Amir Abboud, Virginia Vassilevska Williams, and Huacheng Yu. Matching triangles and basing hardness on an extremely popular conjecture. In *Proceedings of the 47th Annual ACM on Symposium on Theory of Computing, STOC*, pages 41–50, 2015.
- [4] Isolde Adler, Stavros G. Kolliopoulos, Philipp Klaus Krause, Daniel Lokshtanov, Saket Saurabh, and Dimitrios M. Thilikos. Irrelevant vertices for the planar disjoint paths problem. *J. Comb. Theory, Ser. B*, 122:815–843, 2017.
- [5] Jochen Alber and Rolf Niedermeier. Improved tree decomposition based algorithms for domination-like problems. In *Proceedings of the 5th Latin American Symposium on Theoretical Informatics, LATIN*, volume 2286 of *Lecture Notes in Computer Science*, pages 613–628. Springer, 2002.
- [6] Arturs Backurs and Piotr Indyk. Edit distance cannot be computed in strongly sub-quadratic time (unless SETH is false). In *Proceedings of the 47th Annual ACM on Symposium on Theory of Computing, STOC*, pages 51–58, 2015.

- [7] MohammadHossein Bateni, Mohammad Taghi Hajiaghayi, and Dániel Marx. Approximation schemes for steiner forest on planar graphs and graphs of bounded treewidth. *J. ACM*, 58(5):21:1–21:37, 2011.
- [8] Andreas Björklund, Thore Husfeldt, Petteri Kaski, and Mikko Koivisto. Fourier meets Möbius: fast subset convolution. In *Proceedings of the 39th Annual ACM on Symposium on Theory of Computing, STOC*, pages 67–74, 2007.
- [9] Ivan Bliznets, Marek Cygan, Pawel Komosa, Lukás Mach, and Michal Pilipczuk. Lower bounds for the parameterized complexity of minimum fill-in and other completion problems. In *Proceedings of the 27th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1132–1151, 2016.
- [10] Hans L. Bodlaender, Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Deterministic single exponential time algorithms for connectivity problems parameterized by treewidth. *Inf. Comput.*, 243:86–111, 2015.
- [11] Glencora Borradaile and Hung Le. Optimal dynamic program for r-domination problems over tree decompositions. In *Proceedings of the 11th International Symposium on Parameterized and Exact Computation, IPEC*, volume 63 of *LIPICs*, pages 8:1–8:23. Schloss Dagstuhl - Leibniz-Zentrum fuer Informatik, 2016.
- [12] Chris Calabro, Russell Impagliazzo, and Ramamohan Paturi. The complexity of satisfiability of small depth circuits. In *Proceedings of the 4th International Workshop on Parameterized and Exact Computation, IWPEC*, volume 5917 of *Lecture Notes in Computer Science*, pages 75–85. Springer, 2009.
- [13] Shiri Chechik, Daniel H. Larkin, Liam Roditty, Grant Schoenebeck, and Robert Endre Tarjan and Virginia Vassilevska Williams. Better approximation algorithms for the graph diameter. In *Proceedings of the 25th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1041–1052, 2014.
- [14] Jianer Chen, Xiuzhen Huang, Iyad A. Kanj, and Ge Xia. On the computational hardness based on linear FPT-reductions. *J. Comb. Optim.*, 11(2):231–247, 2006.
- [15] Jianer Chen, Xiuzhen Huang, Iyad A. Kanj, and Ge Xia. Strong computational lower bounds via parameterized complexity. *J. Comput. Syst. Sci.*, 72(8):1346–1367, 2006.
- [16] Radu Curticapean and Dániel Marx. Tight conditional lower bounds for counting perfect matchings on graphs of bounded treewidth, cliquewidth, and genus. In *Proceedings of the 27th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1650–1669, 2016.
- [17] Marek Cygan, Holger Dell, Daniel Lokshtanov, Dániel Marx, Jesper Nederlof, Yoshio Okamoto, Ramamohan Paturi, Saket Saurabh, and Magnus Wahlström. On problems as hard as CNF-SAT. *ACM Trans. Algorithms*, 12(3):41:1–41:24, 2016.
- [18] Marek Cygan, Fedor V. Fomin, Alexander Golovnev, Alexander S. Kulikov, Ivan Mihajlin, Jakub Pachocki, and Arkadiusz Socala. Tight bounds for graph homomorphism and subgraph isomorphism. In *Proceedings of the 27th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1643–1649, 2016.
- [19] Marek Cygan, Fedor V. Fomin, Lukasz Kowalik, Daniel Lokshtanov, Dániel Marx, Marcin Pilipczuk, Michal Pilipczuk, and Saket Saurabh. *Parameterized Algorithms*. Springer, 2015.

- [20] Marek Cygan, Stefan Kratsch, and Jesper Nederlof. Fast hamiltonicity checking via bases of perfect matchings. In *Proceedings of the 45th Annual ACM on Symposium on Theory of Computing, STOC*, pages 301–310, 2013.
- [21] Marek Cygan, Jesper Nederlof, Marcin Pilipczuk, Michal Pilipczuk, Johan M. M. van Rooij, and Jakub Onufry Wojtaszczyk. Solving connectivity problems parameterized by treewidth in single exponential time. In *Proceedings of the 52nd Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pages 150–159, 2011.
- [22] Marek Cygan, Marcin Pilipczuk, and Michal Pilipczuk. Known algorithms for Edge Clique Cover are probably optimal. *SIAM J. Comput.*, 45(1):67–83, 2016.
- [23] Erik D. Demaine, Fedor V. Fomin, Mohammad Taghi Hajiaghayi, and Dimitrios M. Thilikos. Subexponential parameterized algorithms on bounded-genus graphs and -minor-free graphs. *J. ACM*, 52(6):866–893, 2005.
- [24] Erik D. Demaine and MohammadTaghi Hajiaghayi. The bidimensionality theory and its algorithmic applications. *Comput. J.*, 51(3):292–302, 2008.
- [25] David Eppstein. Diameter and treewidth in minor-closed graph families. *Algorithmica*, 27(3):275–291, 2000.
- [26] Samuel Fiorini, Nadia Hardy, Bruce A. Reed, and Adrian Vetta. Planar graph bipartization in linear time. *Discrete Applied Mathematics*, 156(7), 2008.
- [27] Jörg Flum and Martin Grohe. *Parameterized Complexity Theory*. Springer, Berlin, 2006.
- [28] Fedor V. Fomin, Serge Gaspers, Saket Saurabh, and Alexey A. Stepanov. On two techniques of combining branching and treewidth. *Algorithmica*, 54(2):181–207, 2009.
- [29] Fedor V. Fomin, Petr A. Golovach, Daniel Lokshtanov, and Saket Saurabh. Almost optimal lower bounds for problems parameterized by clique-width. *SIAM J. Comput.*, 43(5):1541–1563, 2014.
- [30] Fedor V. Fomin, Daniel Lokshtanov, Fahad Panolan, and Saket Saurabh. Efficient computation of representative families with applications in parameterized and exact algorithms. *J. ACM*, 63(4):29:1–29:60, 2016.
- [31] Russell Impagliazzo and Ramamohan Paturi. On the complexity of k -sat. *J. Comput. Syst. Sci.*, 62(2):367–375, 2001.
- [32] Russell Impagliazzo, Ramamohan Paturi, and Francis Zane. Which problems have strongly exponential complexity? *J. Comput. Syst. Sci.*, 63(4):512–530, 2001.
- [33] Lars Jaffke and Bart M. P. Jansen. Fine-grained parameterized complexity analysis of graph coloring problems. In *Proceedings of the 10th International Conference on Algorithms and Complexity, CIAC*, volume 10236 of *Lecture Notes in Computer Science*, pages 345–356, 2017.
- [34] Bart M. P. Jansen, Daniel Lokshtanov, and Saket Saurabh. A near-optimal planarization algorithm. In *Proceedings of the 25th Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1802–1811, 2014.
- [35] Jon Kleinberg and Éva Tardos. *Algorithm Design*. Addison-Wesley Longman Publishing Co., Inc., Boston, MA, USA, 2005.

- [36] Michael Lampis. Parameterized approximation schemes using graph widths. In *Proceedings of the 41st International Colloquium on Automata, Languages, and Programming, ICALP*, volume 8572 of *Lecture Notes in Computer Science*, pages 775–786. Springer, 2014.
- [37] Daniel Lokshtanov, Dániel Marx, and Saket Saurabh. Known algorithms on graphs on bounded treewidth are probably optimal. In *Proceedings of the 22nd Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 777–789, 2011.
- [38] Dániel Marx. On the optimality of planar and geometric approximation schemes. In *Proceedings of the 48th Annual IEEE Symposium on Foundations of Computer Science, FOCS*, pages 338–348, 2007.
- [39] Dániel Marx. Can you beat treewidth? *Theory of Computing*, 6(1):85–112, 2010.
- [40] Dániel Marx and Valia Mitsou. Double-exponential and triple-exponential bounds for choosability problems parameterized by treewidth. In *43rd International Colloquium on Automata, Languages, and Programming, ICALP*, volume 55 of *LIPICs*, pages 28:1–28:15, 2016.
- [41] Daniel Mölle, Stefan Richter, and Peter Rossmanith. Enumerate and expand: Improved algorithms for connected vertex cover and tree cover. *Theory Comput. Syst.*, 43(2):234–253, 2008.
- [42] Jesper Nederlof, Johan M. M. van Rooij, and Thomas C. van Dijk. Inclusion/exclusion meets measure and conquer. *Algorithmica*, 69(3):685–740, 2014.
- [43] Rolf Niedermeier. *Invitation to fixed-parameter algorithms*, volume 31 of *Oxford Lecture Series in Mathematics and its Applications*. Oxford University Press, Oxford, 2006.
- [44] Mihai Patrascu and Ryan Williams. On the possibility of faster SAT algorithms. In *Proceedings of the 21st Annual ACM-SIAM Symposium on Discrete Algorithms, SODA*, pages 1065–1075, 2010.
- [45] Liam Roditty and Virginia Vassilevska Williams. Fast approximation algorithms for the diameter and radius of sparse graphs. In *Proceedings of the 45th Annual ACM on Symposium on Theory of Computing, STOC*, pages 515–524, 2013.
- [46] Alexander D. Scott and Gregory B. Sorkin. Linear-programming design and analysis of fast algorithms for max 2-csp. *Discrete Optimization*, 4(3-4):260–287, 2007.
- [47] Atsushi Takahashi, Shuichi Ueno, and Yoji Kajitani. Mixed searching and proper-path-width. *Theor. Comput. Sci.*, 137(2):253–268, 1995.
- [48] Jan Arne Telle and Andrzej Proskurowski. Practical algorithms on partial k-trees with an application to domination-like problems. In *Proceedings of the 3rd Workshop on Algorithms and Data Structures, WADS*, volume 709 of *Lecture Notes in Computer Science*, pages 610–621. Springer, 1993.
- [49] Dimitrios M. Thilikos, Maria J. Serna, and Hans L. Bodlaender. Cutwidth I: A linear time fixed parameter algorithm. *J. Algorithms*, 56(1):1–24, 2005.
- [50] Johan M. M. van Rooij, Hans L. Bodlaender, and Peter Rossmanith. Dynamic programming on tree decompositions using generalised fast subset convolution. In *Proceedings of the 17th Annual European Symposium on Algorithms, ESA*, volume 5757 of *Lecture Notes in Computer Science*, pages 566–577. Springer, 2009.

- [51] Virginia Vassilevska Williams. Hardness of easy problems: Basing hardness on popular conjectures such as the strong exponential time hypothesis (invited talk). In *Proceedings of the 10th International Symposium on Parameterized and Exact Computation, IPEC*, volume 43 of *LIPICs*, pages 17–29, 2015.