

Geometric graphs with few disjoint edges

Géza Tóth*

Courant Institute, NYU and DIMACS Center, Rutgers University

Pavel Valtr†

Charles University, Prague and DIMACS Center, Rutgers University

Abstract

A *geometric graph* is a graph drawn in the plane so that the vertices are represented by points in general position, the edges are represented by straight line segments connecting the corresponding points.

Improving a result of Pach and Törőcsik, we show that a geometric graph on n vertices with no $k + 1$ pairwise disjoint edges has at most $k^3(n + 1)$ edges. On the other hand, we construct geometric graphs with n vertices and approximately $\frac{3}{2}(k - 1)n$ edges, containing no $k + 1$ pairwise disjoint edges.

We also improve both the lower and upper bounds of Goddard, Katchalski and Kleitman on the maximum number of edges in a geometric graph with no four pairwise disjoint edges.

1 Introduction

A *geometric graph* G is a graph drawn in the plane by (possibly crossing) straight line segments, i.e., it is defined as a pair $G = (V, E)$, where V is a set of points in general position in the plane and E is a set of closed segments whose endpoints belong to V .

The following problem was raised by Avital and Hanani [AH], Kupitz [K], Erdős and Perles. Determine the smallest number $e_k(n)$ such that any geometric graph with n vertices and $m > e_k(n)$ edges contains $k + 1$ pairwise disjoint edges. By a result of Hopf and Pannwitz [HP] and Erdős, $e_1(n) = n$. Alon and Erdős [AE] showed that $e_2(n) \leq 6n - 5$ which was improved by Goddard, Katchalski and Kleitman [GKK] to $e_2(n) \leq 3n$. The best known lower bound, $e_2(n) \geq 2.5n - 4$, is due to Perles (see [PA], [MP]). It is also known that $3.5n - 10 \leq e_3(n) \leq 10n$ [GKK]. For any fixed k , Pach and Törőcsik [PT] were first to prove that $e_k(n)$ is linear in n ;

*Supported by OTKA-T-020914, OTKA-F-22234 and NSF grant CCR-94-24398.

†Supported by the Czech Republic grant GAČR 201/1996/0194 and Charles University grants GAUK 193, 194/1996

their bound was $e_k(n) \leq k^4 n$. It follows from a result of Kupitz [K] that $e_k(n) \geq kn$. In this paper we further improve both the upper and lower bounds for general k .

Theorem 1 For $k \leq n/2$,

$$\frac{3}{2}(k-1)n - 2k^2 \leq e_k(n) \leq k^3(n+1).$$

We also improve the above mentioned bounds on $e_3(n)$.

Theorem 2 For any $n \geq 6$,

$$4n - 9 \leq e_3(n) \leq 8.5n.$$

Theorem 1 and Theorem 2 are proved in Sections 2 and 3, respectively. Throughout the paper, we do not make any notational distinction between an edge and the segment representing it.

Recently, using similar methods, the first named author has further improved the upper bound in Theorem 1 to $e_k \leq 2^9 k^2 n$ [T]. For the corresponding problem concerning pairwise *crossing* edges, see [V] and for another related problem see [LMPT].

2 The general case

2.1 The upper bound

Our proof, as the proof of Pach and Töröcsik [PT], is based on Dilworth's theorem [D].

Dilworth's Theorem. *Let P be a partially ordered set containing no chain (totally ordered subset) of size $k+1$. Then P can be covered by k antichains (subsets of pairwise incomparable elements).*

Let $G = (V, E)$ be a geometric graph on n vertices, containing no $k+1$ pairwise disjoint edges. For a vertex v , let $x(v)$ and $y(v)$ denote its x - and y -coordinate, respectively. We can assume without loss of generality that no two vertices have the same x -coordinate.

An edge e is said to *lie below* an edge e' , if no vertical line crossing both e and e' crosses e strictly above e' . Finally, let $\pi(e)$ denote the orthogonal projection of e to the x -axis.

Define four binary relations \prec_i ($i = 1, \dots, 4$) on the edge set E as follows (see also [PT, PA]). Let $e = v_1 v_2, e' = v'_1 v'_2$ be two disjoint edges of G , where $x(v_1) < x(v_2)$ and $x(v'_1) < x(v'_2)$. Then (see Fig. 1)

$$\begin{aligned} e \prec_1 e', & \text{ if } x(v_1) < x(v'_1), x(v_2) < x(v'_2), \text{ and } e \text{ lies below } e', \\ e \prec_2 e', & \text{ if } x(v_1) > x(v'_1), x(v_2) > x(v'_2), \text{ and } e \text{ lies below } e', \end{aligned}$$




Figure 1: The relations \prec_i .

$$\begin{aligned}
 e \prec_3 e', & \text{ if } x(v_1) < x(v'_1), x(v_2) > x(v'_2), \text{ and } e \text{ lies below } e', \\
 e \prec_4 e', & \text{ if } x(v_1) > x(v'_1), x(v_2) < x(v'_2), \text{ and } e \text{ lies below } e'.
 \end{aligned}$$

Obviously, each of the relations \prec_i is a partial ordering, and any pair of disjoint edges in G is comparable by at least one of them.

Since G does not contain $k+1$ disjoint edges, (E, \prec_1) does not contain a chain of length $k+1$. Therefore, by Dilworth's theorem, E can be covered by k antichains with respect to \prec_1 . Let E_1 be the largest of these antichains, thus $|E_1| \geq |E|/k$. Applying Dilworth's theorem on (E_1, \prec_2) , we similarly get an antichain (with respect to \prec_2) $E_2 \subseteq E_1$ of size $|E_2| \geq |E_1|/k \geq |E|/k^2$. In the rest of the proof we estimate the size of E_2 from above.

Since E_2 is an antichain with respect to \prec_1 and \prec_2 , $\pi(e) \cap \pi(e') \neq \emptyset$ for any $e, e' \in E_2$. Therefore, $\bigcap_{e \in E_2} \pi(e) \neq \emptyset$, so there is a vertical line ℓ which intersects all edges in E_2 .

Let $\vec{G}_2 = (V, \vec{E}_2)$ be a directed geometric graph obtained from (V, E_2) by replacing each

edge $e = v_1v_2$ in E_2 by the two oriented edges $\overrightarrow{v_1v_2}$ and $\overrightarrow{v_2v_1}$. For two edges $e_1 = \overrightarrow{v_0v_1}$, $e_2 = \overrightarrow{v_1v_2}$ forming a path in $\overrightarrow{G_2}$, we say that e_2 is a *zag* of e_1 , if the following two conditions hold:

- (i) $\pi(e_1) \cap \pi(e_2)$ has positive length,
- (ii) for any $z \in (\pi(e_1) \cap \pi(e_2)) \setminus \{v_1\}$, the vertical line through z intersects e_2 below e_1 , and it intersects no other edge going from v_1 between e_1 and e_2 .

Observe that each edge in $\overrightarrow{G_2}$ has at most one zag. We call an oriented path $e_1e_2 \dots e_r$ in $\overrightarrow{G_2}$ a *zigzag path*, if e_{i+1} is the zag of e_i , for each $i = 1, \dots, r - 1$ (see Fig. 2).




Figure 2: Maximal zigzag paths.

Lemma 3 *Every zigzag path in $\overrightarrow{G_2}$ has at most $2k$ edges.*

Lemma 4 *$\overrightarrow{G_2}$ has at most $n + 1$ maximal zigzag paths.*

Lemmas 3 and 4 immediately give Theorem 1: every edge in $\overrightarrow{G_2}$ is contained in a maximal zigzag path; therefore, by Lemmas 3 and 4,

$$|\overrightarrow{E_2}| \leq 2k(n + 1),$$

and, consequently,

$$|E| \leq k^2 |E_2| = k^2 |\vec{E}_2|/2 \leq k^3(n+1).$$

It remains to prove Lemmas 3 and 4.

Proof of Lemma 3. Let $P = e_1 e_2 \dots e_r$ be a zigzag path in \vec{G}_2 , and let $e_i = \overrightarrow{v_{i-1} v_i}$ for $i = 1, \dots, r$. We need the following claim.

Claim One of the two sequences

$$S_0 = x(v_0), x(v_2), \dots \quad S_1 = x(v_1), x(v_3), \dots$$

is decreasing, and the other one is increasing.

Proof of Claim. Suppose the claim is false. Then there is an index i such that either $x(v_{i-1}) < x(v_{i+1})$ and $x(v_i) < x(v_{i+2})$, or $x(v_{i-1}) > x(v_{i+1})$ and $x(v_i) > x(v_{i+2})$. Consequently, $e_{i+2} \prec_1 e_i$ or $e_{i+2} \prec_2 e_i$ (see Fig. 3 for all four possible cases), which is a contradiction




Figure 3: $e_{i+2} \prec_1 e_i$ or $e_{i+2} \prec_2 e_i$.

with the definition of E_2 . The claim is proved. \square

The length of P is at most $2k$ since otherwise $e_1, e_3, \dots, e_{2k+1}$ would be $k+1$ disjoint edges according to Claim (see Fig. 4). \square

Proof of Lemma 4. Let $v \in V$ be a vertex lying to the right of ℓ , and let P_1, P_2 be two different zigzag paths in \vec{G}_2 ending in v . If the slope of the last edge in P_1 is, say, smaller than the slope of the last edge in P_2 , then the last edge of P_1 has a zag and, consequently, P_1 can be extended to a longer zigzag path. It follows that at most one maximal zigzag path ends in v , and this is similarly true for any vertex not lying on ℓ . Similarly, at most two zigzag paths end in any vertex v on ℓ (one coming to v from the left, the other one from the right). The lemma now follows from the assumption that no pair of vertices lies on a vertical line. \square

2.2 The lower bound

For simplicity, suppose that k is even and n is odd. Set $z = (n - k + 1)/2$. Let P be a set of z points p_1, \dots, p_z placed equidistantly in this order from left to right on a horizontal line p .




Figure 4: $e_1, e_3, \dots, e_{2k+1}$ are disjoint.

Let $Q = \{q_1, \dots, q_z\}$ be a translation of P such that q_i always corresponds to p_i and $p_1 p_z q_z q_1$ is a square. Let q be the line containing Q , and let r be the line parallel to p and q , halfway between them. Let E_1 be the set of all segments (edges) $p_i q_j$ with $-k/2 \leq i + j - (z + 1) \leq k/2$ (see Fig. 5).

The segments of E_1 intersect the line r in $k + 1$ points $r_{-k/2}, r_{-k/2+1}, \dots, r_{k/2}$ such that r_t always lies on the segments $p_i q_j$ with $i + j - (z + 1) = t$. Let R be the set of the centers of the segments $r_t r_{t+1}$ ($t = -k/2, -k/2 + 1, \dots, k/2 - 2$), and let E_2 be the set of all edges (segments) joining vertices of R with vertices of $P \cup Q$ (see Fig. 5).

We now show that the geometric graph $G = (P \cup Q \cup R, E_1 \cup E_2)$ gives the lower bound in Theorem 1.

First, observe that G has $z + z + (k - 1) = n$ vertices and

$$\begin{aligned} |E_1| + |E_2| &= \frac{(n - k + 1)(k + 1) - 4(1 + 2 + \dots + k/2)}{2} + (n - k + 1)(k - 1) \\ &= (n - k + 1)\left(\frac{3k}{2} - \frac{1}{2}\right) - \frac{k(k + 2)}{4} > \frac{3}{2}(k - 1)n - 2k^2 \end{aligned}$$

edges. It remains to show that G contains no $k + 1$ pairwise disjoint edges. Suppose that D is a set of disjoint edges in G . If $|D \cap E_1| \leq 1$, then $|D| \leq |R| + 1 = k$. Otherwise, let the leftmost edge of $D \cap E_1$, e_1 , intersect r in a point r_s , and the rightmost one, e_2 , in a point r_t . Since there are $s + k/2$ vertices of R to the left of e_1 , D contains at most $s + k/2$ edges to the left of e_1 . Similarly, D contains at most $k/2 - t$ edges to the right of e_2 . Since there are $t - s - 2$ vertices of $P \cup Q$ between e_1 and e_2 , D contains at most $t - s - 2$ edges between e_1 and e_2 . Altogether, D contains at most $(s + k/2) + (k/2 - t) + (t - s - 2) + 2 = k$ edges. \square




Figure 5: The edges of E_1 and E_2 (for $k = 4$, $n = 15$, $z = 6$).

3 The case $k = 3$

3.1 The upper bound

We use the following result.

Lemma 5 ([GKK]) *If a geometric graph G of n vertices does not contain 4 pairwise disjoint edges and there is a line which intersects every edge of G and contains no vertex of G , then G has at most $7n$ edges.*

Lemma 5 is not stated in [GKK] explicitly. However, its proof (relatively long case-analysis) is readily contained in the proof of Theorem 2 in [GKK].

Let $G = (V, E)$ be a geometric graph without 4 pairwise disjoint edges. Denote the vertices by v_1, \dots, v_n from left to right, and assume that no pair of them lies on a vertical line. For any $1 \leq i < n$ let \overline{G}_i be the subgraph of G which contains only those edges $v_\alpha v_\beta$ of G where $\alpha \leq i < \beta$.

It follows from Lemma 5 and the assumption that for any $1 \leq i < n$ $|E(\overline{G}_i)| \leq 7n$. For any $1 \leq i < n$, let G_i^- and G_i^+ be the subgraph of G induced by the vertices v_1, \dots, v_i and by v_i, \dots, v_n , respectively. Let

$$I = \max \left\{ i \mid G_i^- \text{ does not contain 2 disjoint edges} \right\}.$$

Since G_{I+1}^- contains two disjoint edges, G_{I+2}^+ does not. Suppose without loss of generality that $I < n/2$. Then

$$E(G) = E(G_I^-) \cup E(G_{I+2}^+) \cup E(\overline{G}_{I+1}) \cup \{v_i v_{I+1} \in E(G) \mid i \leq I\}.$$

Therefore,

$$|E(G)| \leq e_1(I) + e_1(n - I - 1) + 7n + I < 8.5n.$$

□

3.2 The lower bound

First, let n be odd. Take the vertices of a regular $(n - 2)$ -gon, and join each of them with the furthestmost 4 vertices. Add two vertices near the center of the $(n - 2)$ -gon, and join each of them by an edge to all the other vertices. The resulting geometric graph on n vertices has $4(n - 2)/2 + 2(n - 1) - 1 = 4n - 7$ edges. It is easy to see that no 4 edges are pairwise disjoint, see Fig. 6.




Figure 6: A geometric graph with no 4 pairwise disjoint edges.

For n even, we take the above geometric graph on $n + 1$ vertices, and remove a vertex and the six edges incident to it. The resulting graph on n vertices has $4n - 9$ edges. □

4 Remarks

By a little modification, it is possible to slightly improve the upper bound of Theorem 1.

Theorem 6 For any $k \leq n/2$,

$$e_k(n) \leq \frac{4}{21}k^3n + O(k^2n).$$

Proof. (Sketch) We use the same partial orderings, $\prec_1, \prec_2, \prec_3, \prec_4$, as in the proof of Theorem 1 (see Fig. 1). For any edge e of a geometric graph $G = (V, E)$, let $\text{Rank}_3^+(e)$ be the maximum number $a \geq 0$ such that there exist $e_1, e_2, \dots, e_a \in E$, $e \prec_3 e_1 \prec_3 e_2 \prec_3 e_3 \cdots \prec_3 e_a$, and let $\text{Rank}_4^-(e)$ be the maximum number $b \geq 0$ such that there exist $e_1, e_2, \dots, e_b \in E$, $e_1 \prec_4 e_2 \prec_4 e_3 \cdots \prec_4 e_b \prec_4 e$.

Lemma 7 Let $G = (V, E)$ be a geometric graph with n vertices and with no $k + 1$ pairwise disjoint edges. Suppose that there is a line which avoids all vertices and crosses all edges of G . Then

$$|E| \leq \frac{1}{6}k^3n + O(k^2n).$$

Proof of Lemma 7. For any $0 \leq a, b \leq k - 1$ let

$$E_{a,b} = \left\{ e \in E \mid \text{Rank}_3^+(e) = a, \text{Rank}_4^-(e) = b \right\}.$$

Clearly, $E_{a,b}$ is an antichain with respect to \prec_3 and \prec_4 . We can define $\overrightarrow{E_{a,b}}$ and its maximal zigzag paths just as in the proof Theorem 1. It is easy to see that there are at most n maximal zigzag paths and every edge of $\overrightarrow{E_{a,b}}$ is contained in at least one of them (in fact, in exactly one of them, but here we do not need that). Suppose that there is a zigzag path $\overrightarrow{e_1}, \overrightarrow{e_2}, \dots, \overrightarrow{e_x}$ of $x = 2(k - a - b) + 1$ edges. Then $\overrightarrow{e_1}, \overrightarrow{e_3}, \overrightarrow{e_5}, \dots, \overrightarrow{e_x}$ are $k - a - b + 1$ pairwise disjoint edges. Moreover, we can add a edges above $\overrightarrow{e_1}$ and b edges below $\overrightarrow{e_x}$ to get $k + 1$ pairwise disjoint edges, contradicting our assumption. (See Fig. 7.)

Therefore, every maximal zigzag path of $\overrightarrow{E_{a,b}}$ has at most $2(k - a - b)$ edges, so

$$|E_{a,b}| \leq n(k - a - b),$$

$$|E| = \sum_{\substack{0 \leq a, b \\ a+b < k}} |E_{a,b}| \leq n \sum_{\substack{0 \leq a, b \\ a+b < k}} (k - a - b) = \binom{k+2}{3} n.$$

□




Figure 7: $k + 1$ disjoint edges.

Return to the proof of Theorem 6. It is not hard to see that there is a line ℓ which avoids all vertices of G and on one side of ℓ there are $\lceil n/2 \rceil$ vertices and at most $\lceil k/2 \rceil$ pairwise disjoint edges, while on the other side of ℓ there are $\lfloor n/2 \rfloor$ vertices and at most $\lfloor k/2 \rfloor$ pairwise disjoint edges of G . We get the recursion

$$e_k(n) \leq \frac{1}{6}k^3n + O(k^2)n + e_{\lceil k/2 \rceil}(\lceil n/2 \rceil) + e_{\lfloor k/2 \rfloor}(\lfloor n/2 \rfloor)$$

and Theorem 6 follows. \square

If the number of edges in a geometric graph is at least $\Omega(n^2)$, then Theorem 1 guarantees $\Omega(n^{1/3})$ pairwise disjoint edges. This is improved by the following result of Pach [P].

Theorem 8 ([P]) *For any $c > 0$ there is a $c' > 0$ such that every geometric graph of n vertices and at least cn^2 edges has at least $c'\sqrt{n}$ pairwise disjoint edges.*

References

[AE] N. Alon and P. Erdős, Disjoint edges in geometric graphs, *Discrete and Computational Geometry* 4 (1989), 287-290.

- [AH] S. Avital and H. Hanani, Graphs, *Gilyonot Lematematika* **3** (1966), 2-8 (in Hebrew).
- [D] R. P. Dilworth, A decomposition theorem for partially ordered sets, *Annals of Mathematics* **51** (1950), 161-166.
- [GKK] W. Goddard, M. Katchalski, and D. J. Kleitman, Forcing disjoint segments in the plane, *European Journal of Combinatorics* **17** (1996), 391-395.
- [HP] H. Hopf and E. Pannwitz, Aufgabe No. 167, *Jahresbericht der Deutschen Mathematiker-Vereinigung* **43** (1934), 114.
- [K] Y. Kupitz, Extremal problems in combinatorial geometry, *Aarhus University Lecture Notes Series*, **53**, (1979), Aarhus University, Denmark.
- [LMP] D. Larman, J. Matoušek, J. Pach, and J. Töröcsik, A Ramsey-type result for planar convex sets, *The Bulletin of the London Mathematical Society* **26**, (1994), 132-136.
- [MP] W. Moser and J. Pach, Recent developments in combinatorial geometry, in: *New trends in discrete and computational geometry (J. Pach, ed.)*, Algorithms and Combinatorics **10**, Springer, Berlin, 1993, 281-302.
- [P] J. Pach, personal communication.
- [PA] J. Pach and P.K. Agarwal, *Combinatorial Geometry*, John Wiley, New York, 1995.
- [PT] J. Pach and J. Töröcsik, Some geometric applications of Dilworth's theorem, *Discrete and Computational Geometry* **12** (1994), 1-7.
- [T] G. Tóth, Note on geometric graphs, (manuscript)
- [V] P. Valtr, On geometric graphs with no k pairwise parallel edges, *Discrete and Computational Geometry* **19** (1998) 461-469.